arXiv:0812.3343v6 [math.QA] 21 Jun 2009

Convex PBW-Type Lyndon Bases and Restricted Two-parameter Quantum Groups of Type B

Naihong Hu[⋆] and Xiuling Wang[*]

Abstract. We construct convex PBW-type Lyndon bases for two-parameter quantum groups $U_{r,s}(\mathfrak{so}_{2n+1})$ with detailed commutation relations. It turns out that under a certain condition, the restricted two-parameter quantum group $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ (r, s are roots of unity) is of Drinfeld double. All of Hopf isomorphisms of $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$, as well as $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ are determined. Finally, necessary and sufficient conditions for $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ to be a ribbon Hopf algebra are singled out by describing the left and right integrals.

1. Introduction

1.1. In 2001, from the down-up algebras approach, Benkart-Witherspoon in [**BW1**] reobtained Takeuchi's definition of two-parameter quantum groups of type A. Since then, a systematic study of the two-parameter quantum groups has been going on, for instance, [**BW2, BW3**] for type A; [**BGH1, BGH2**] for type B, C, D; [**HS, BH**] for types G_2, E_6, E_7, E_8. For a unified definition, see [**HP**]. Hu-Rosso-Zhang [**HRZ**] defined the affine type A case, obtained its Drinfeld realization in two-parameter quantum version and constructed the quantum affine Lyndon basis. Furthermore, the vertex operator representations of level 1 for the two-parameter quantum affine algebras $U_{r,s}(\widehat{\mathfrak{g}})$ can be established (cf. [**HZ**], etc.). While, in the root of unity case, Hu-Wang [**HW**] generalized the work [**BW3**] to the restricted type G_2 case. The goal of this paper is to solve the restricted type B case with overcoming some peculiar difficulties itself in this case.

1.2. In the study of Lusztig's symmetry property for the two-parameter quantum groups in question, Theorem 3.1 [**BGH1**] says that the Lusztig's symmetries only exist as $\mathbb{Q}$-isomorphisms between $U_{r,s}(\mathfrak{g})$ and its associated object $U_{s^{-1},r^{-1}}(\mathfrak{g})$ when $\text{rank}(\mathfrak{g}) = 2$, and in the case when $\text{rank}(\mathfrak{g}) > 2$, the "*iff*" condition for the existence of Lusztig's symmetries forces $U_{r,s}(\mathfrak{g})$ to take the "one-parameter" form $U_{q,q^{-1}}(\mathfrak{g})$, where $r = q = s^{-1}$. This means that one cannot conveniently write out the convex

---

1991 *Mathematics Subject Classification.* Primary 17B37, 81R50; Secondary 17B35.

Key words and phrases. Convex PBW-type Lyndon basis, restricted 2-parameter quantum groups, integrals, ribbon Hopf algebra.

[⋆] N.H., Corresponding Author, supported in part by the NNSF (Grant 10728102), the PCSIRT and the from the MOE, the National & Shanghai Leading Academic Discipline Projects (Project Number: B407).

[*] X.L., supported by the Nankai Research-encouraging Fund for the PhD-Teachers, and a fund from LPMC.

PBW-type basis for $U_{r,s}(\mathfrak{g})$ like in the one-parameter cases using Lusztig's braid group actions in a typical fashion (see [**Lu**]). This is one of difficulties encountered in the two-parameter setting. Note that the combinatorial construction of the PBW-type bases in the quantum cases is available but nontrivial, as it depends not only on the choice of a convex ordering ([**B, Ro**]) on a positive root system, but also the inserting manner of the $\mathbf{q}$-bracketings with suitable structure constants (see [**Ro, HRZ**], etc.) into the good Lyndon words (cf. [**LR, Ro**]). The constructions of convex PBW-type bases in the two-parameter cases have been given for type A in [**BW3**], types E_6, E_7, E_8 in [**BH**], for types B_2, C_2, D_4 in [**H**], and for type G_2 in [**HW**]. Indeed, in order to study the properties of restriced quantum groups as finite-dimensional Hopf algebras, it is necessary to find the nice bases for the corresponding nonrestricted quantum groups at generic cases. The first thing of the article is to present a direct construction for a convex PBW-type Lyndon basis of type B (for arbitrary rank) and provide explicit information on commutation relations among basis elements that is decisive to single out the left (or right) integrals of $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$. Note that the complexity of Lyndon bases corresponding to the nonsimply-laced Dynkin diagrams causes more inconvenience when treated with the type B case here.

1.3. The paper is organized as follows. Section 2 is to give the construction of Lyndon bases for $U_{r,s}(\mathfrak{so}_{2n+1})$. In Section 3, we contribute more efforts to make the possible commutation relations clearly. These give rise to central elements of degree ℓ in the case when parameters r, s are roots of unity, which generate a Hopf ideal. The quotient is the restricted two-parameter quantum groups $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ in Section 4. In Section 5, we show that these Hopf algebras are pointed, and determine all Hopf isomorphisms of $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$, as well as $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ in terms of the set of (left) right skew-primitive elements. Section 6 is to prove that $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ is of Drinfel'd double under a certain condition. In Section 7, the left and right integrals of $\mathfrak{b}$ are singled out based on the discussion in Section 3, which are applied to determine the necessary and sufficient conditions for $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ to be ribbon in Section 8.

2. $U_{r,s}(\mathfrak{so}_{2n+1})$ and its convex PBW-type Lyndon basis

2.1. Let $\mathbb{K} = \mathbb{Q}(r, s)$ be a subfield of $\mathbb{C}$, where $r, s \in \mathbb{C}^*$ with assumptions $r^3 \neq s^3$ and $r^4 \neq s^4$. Let Φ be a root system of $\mathfrak{so}_{2n+1}$ with Π a base of simple roots, which is a finite subset of a Euclidean space $E = \mathbb{R}^3$ with an inner product $(\ ,\ )$. Let $\epsilon_1, \cdots, \epsilon_n$ denote an orthonormal basis of E, then $\Pi = \{\alpha_i = \epsilon_i - \epsilon_{i+1} \mid 1 \leq i < n\} \cup \{\alpha_n = \epsilon_n\}$, $\Phi = \{\pm\epsilon_i \pm \epsilon_j \mid 1 \leq i \neq j \leq n\} \cup \{\pm\epsilon_i \mid 1 \leq i \leq n\}$. In this case, set $r_i = r^{(\alpha_i,\, \alpha_i)}$, $s_i = s^{(\alpha_i,\, \alpha_i)}$, so that $r_1 = \cdots = r_{n-1} = r^2, r_n = r$ and $s_1 = \cdots = s_{n-1} = s^2, s_n = s$.

Given two sets of symbols $W = \{\omega_1, \cdots, \omega_n\}$, $W' = \{\omega_1', \cdots, \omega_n'\}$. Define the structural constants matrix $(\langle\omega_i', \omega_j\rangle)_{n\times n}$ of type B by

$$\begin{pmatrix} r^2s^{-2} & r^{-2} & 1 & \cdots & 1 & 1 \\ s^2 & r^2s^{-2} & r^{-2} & \cdots & 1 & 1 \\ 1 & s^2 & r^2s^{-2} & \cdots & 1 & 1 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & 1 & 1 & \cdots & r^2s^{-2} & r^{-2} \\ 1 & 1 & 1 & \cdots & s^2 & rs^{-1} \end{pmatrix}.$$

Definition 2.1. ([**BGH1**]) Let $U = U_{r,s}(\mathfrak{so}_{2n+1})$ be the associative algebra over $\mathbb{K}$ generated by $e_i,\ f_i,\ \omega_i^{\pm 1},\ \omega_i'^{\pm 1}$ $(1 \le i \le n)$ subject to relations $(B1)-(B5)$:

$(B1)$ The $\omega_i^{\pm1}, \omega_j'^{\pm1}$ all commute with one another and $\omega_i\omega_i^{-1} = 1 = \omega_j'\omega_j'^{-1}$.

$$(B2) \qquad \omega_j\, e_i\, \omega_j^{-1} = \langle \omega_i', \omega_j\rangle e_i, \qquad \omega_j\, f_i\, \omega_j^{-1} = \langle \omega_i', \omega_j\rangle^{-1} f_i.$$

$$(B3) \qquad \omega_j'\, e_i\, \omega_j'^{-1} = \langle \omega_j', \omega_i\rangle^{-1} e_i, \qquad \omega_j'\, f_i\, \omega_j'^{-1} = \langle \omega_j', \omega_i\rangle f_i.$$

$$(B4) \qquad [\, e_i, f_j\,] = \delta_{ij}\frac{\omega_i - \omega_i'}{r_i - s_i}.$$

$(B5)$ (r,s)-Serre relations

$$(\mathrm{ad}_\ell e_i)^{1-a_{ij}}(e_j) = 0, \qquad (\mathrm{ad}_r f_i)^{1-a_{ij}}(f_j) = 0$$

are given by its Hopf algebra structure on $U_{r,s}(\mathfrak{so}_{2n+1})$ with the comultiplication, the counit and the antipode as follows

$$\begin{gathered}
\Delta(\omega_i^{\pm1}) = \omega_i^{\pm1}\otimes\omega_i^{\pm1}, \qquad \Delta(\omega_i'^{\pm1}) = \omega_i'^{\pm1}\otimes\omega_i'^{\pm1},\\
\Delta(e_i) = e_i\otimes 1 + \omega_i\otimes e_i, \qquad \Delta(f_i) = 1\otimes f_i + f_i\otimes\omega_i',\\
\varepsilon(\omega_i^{\pm}) = \varepsilon(\omega_i'^{\pm1}) = 1, \qquad \varepsilon(e_i) = \varepsilon(f_i) = 0,\\
S(\omega_i^{\pm1}) = \omega_i^{\mp1}, \qquad S(\omega_i'^{\pm1}) = \omega_i'^{\mp1},\\
S(e_i) = -\omega_i^{-1}e_i, \qquad S(f_i) = -f_i\,\omega_i'^{-1}.
\end{gathered}$$

When $r = q = s^{-1}$, $U_{r,s}(\mathfrak{so}_{2n+1})/I \cong U_q(\mathfrak{so}_{2n+1})$, the standard Drinfel'd-Jimbo quantum group, where $I = (\omega_i' - \omega_i^{-1}, \forall\, i)$ is a Hopf ideal.

2.2. U has a triangular decomposition $U \cong U^-\otimes U^0\otimes U^+$, where U^0 is the subalgebra generated by $\omega_i^{\pm1}, \omega_i'^{\pm1}$, and U^+ (resp. U^-) is the subalgebra generated by e_i (resp. f_i). Let $\mathcal{B}$ (resp. $\mathcal{B}'$) denote the Hopf subalgebra of U generated by $e_j, \omega_j^{\pm1}$ (resp. $f_j, \omega_j'^{\pm1}$) with $1 \le j \le n$.

Proposition 2.2. ([**BGH1**]) *There exists a unique skew-dual pairing* $\langle\,,\rangle : \mathcal{B}'\times\mathcal{B}\to\mathbb{K}$ *of the Hopf subalgebras* $\mathcal{B}$ *and* $\mathcal{B}'$ *such that*

$$\begin{gathered}
\langle f_i,\, e_j\rangle = \delta_{ij}\frac{1}{s_i - r_i}, \qquad 1\le i,\, j\le n,\\
\langle {\omega'}_i^{\pm1}, \omega_j^{-1}\rangle = \langle {\omega'}_i^{\pm1}, \omega_j\rangle^{-1} = \langle {\omega'}_i, \omega_j\rangle^{\mp1}, \quad 1\le i,\, j\le n
\end{gathered}$$

and all other pairs of generators are 0. *Moreover, we have* $\langle S(a),\, S(b)\rangle = \langle a,\, b\rangle$ *for* $a\in\mathcal{B}', b\in\mathcal{B}$.

2.3. Write the positive root system $\Phi^+ = \{\alpha_{ij},\, \epsilon_\ell,\, \beta_{ij} \mid 1\le i<j\le n, 1\le \ell\le n\}$, where $\alpha_{ij} := \alpha_i + \alpha_{i+1} + \cdots + \alpha_{j-1} = \epsilon_i - \epsilon_j$, $\epsilon_\ell = \alpha_\ell + \alpha_{\ell+1} + \cdots + \alpha_n$, $\beta_{ij} := \alpha_i + \alpha_{i+1} + \cdots + 2\alpha_n + \alpha_{n-1} + \cdots + \alpha_j = \epsilon_i + \epsilon_j$. To a reduced expression of the maximal length element of Weyl group W for type B_n (see [**CX**]) taken as

$$w_0 = (s_1s_2\cdots s_{n-1}s_ns_{n-1}\cdots s_2s_1)\cdots(s_{n-1}s_ns_{n-1})(s_n),$$

there corresponds a convex ordering on Φ^+ below:

$$\begin{array}{ccccccccc}
\alpha_{12}, & \alpha_{13}, & \cdots, & \alpha_{1n}, & \epsilon_1, & \beta_{1n}, & \cdots, & \beta_{13}, & \beta_{12}, \\
 & \alpha_{23}, & \cdots, & \alpha_{2n}, & \epsilon_2, & \beta_{2n}, & \cdots, & \beta_{23}, & \\
 & & & & \cdots & & & & \\
 & & & \alpha_{n-1,n}, & \epsilon_{n-1}, & \beta_{n-1,n}, & & & \\
 & & & & \epsilon_n. & & & &
\end{array}$$

Remark: The above ordering also corresponds to the standard Lyndon tree of type B_n ([**LR**]):



Denote by $\mathcal{E}_\alpha$ the quantum root vector in U^+ for each $\alpha \in \Phi^+$. Briefly, we set $\mathcal{E}_{i,j} := \mathcal{E}_{\alpha_{i,j+1}}$ for $1 \le i \le j < n$; $\mathcal{E}_{i,n} := \mathcal{E}_{\epsilon_i}$ for $1 \le i \le n$; $\mathcal{E}_{i,j'} := \mathcal{E}_{\beta_{i,j}}$ for $1 \le i < j \le n$; in particular, $\mathcal{E}_{i,i} := \mathcal{E}_{\alpha_i} = e_i$ for $1 \le i \le n$, where in our notation $\alpha_i = \alpha_{i,i+1}$ $(i < n)$ and $\alpha_n = \epsilon_n$.

2.4. With the above ordering and notation, following the grammatical rule of good Lyndon words with inserting (r,s)-bracketings, we can make an inductive definition starting from the bottom to up of the above root-ordering graph as follows.

Definition 2.3. For $\alpha \in \Phi^+$, define $\mathcal{E}_\alpha$ inductively

$$\mathcal{E}_{i,i} = e_i, \qquad 1 \le i \le n, \tag{2.1}$$

$$\mathcal{E}_{i,j} = e_i\mathcal{E}_{i+1,j} - r^2\mathcal{E}_{i+1,j}e_i, \qquad 1 \le i < j \le n, \tag{2.2}$$

$$\mathcal{E}_{i,n'} = \mathcal{E}_{i,n}e_n - rse_n\mathcal{E}_{i,n}, \qquad 1 \le i \le n-1, \tag{2.3}$$

$$\mathcal{E}_{i,j'} = \mathcal{E}_{i,j+1'}e_j - s^{-2}e_j\mathcal{E}_{i,j+1'}, \qquad 1 \le i < j \le n-1. \tag{2.4}$$

Then the (r,s)-Serre relations for U^+ in $(B5)$ can be reformulated as

$$e_i\mathcal{E}_{i,i+1} = s^2\mathcal{E}_{i,i+1}e_i, \tag{2.5}$$

$$\mathcal{E}_{j,j+1}e_{j+1} = s^2e_{j+1}\mathcal{E}_{j,j+1}, \tag{2.6}$$

$$\mathcal{E}_{n-1,n'}e_n = s^2e_n\mathcal{E}_{n-1,n'}, \tag{2.7}$$

for $1 \le i \le n-1$, $1 \le j < n-1$.

Remark: (i) From [**Ro**], the set of quantum Lie brackets (or say, **q**-bracketings) of all good Lyndon words consists of all quantum root vectors of U^+ (for definitions to see *Remark* 3.7 (3) in [**HRZ**], pp. 467).

(ii) As for how to insert (r,s)-bracketings into each good Lyndon word in type B case, in [**H**] an observation was obtained via a representation theoretic analysis in rank 2 case, that is, we have to obey the defining rule as $\mathcal{E}_\gamma := [\mathcal{E}_\alpha, \mathcal{E}_\beta]_{\langle\omega'_\alpha,\omega_\beta\rangle^{-1}} = \mathcal{E}_\alpha\mathcal{E}_\beta - \langle\omega'_\alpha,\omega_\beta\rangle^{-1}\mathcal{E}_\beta\mathcal{E}_\alpha$ for $\alpha, \gamma, \beta \in \Phi^+$ such that $\alpha < \gamma < \beta$ in the convex ordering, and $\gamma = \alpha + \beta$.

2.5. According to [**H, K, Ro, LR**], we have

Theorem 2.4. $\Big\{\mathcal{E}_{n,n}^{c_{n^2}}\cdot\mathcal{E}_{n-1,n'}^{c_{n^2-1}}\mathcal{E}_{n-1,n}^{c_{n^2-2}}\mathcal{E}_{n-1,n-1}^{c_{n^2-3}}\cdots\mathcal{E}_{2,3'}^{c_{4n-4}}\cdots\mathcal{E}_{2,n'}^{c_{3n-1}}\mathcal{E}_{2,n}^{c_{3n-2}}\cdots\mathcal{E}_{2,2}^{c_{2n}}$ $\mathcal{E}_{1,2'}^{c_{2n-1}}\cdots\mathcal{E}_{1,n'}^{c_{n+1}}\mathcal{E}_{1,n}^{c_n}\cdots\mathcal{E}_{1,2}^{c_2}\mathcal{E}_{1,1}^{c_1}\;\Big|\;(c_1,\cdots,c_{n^2}) \in \mathbb{Z}_+^{n^2}\Big\}$ *forms a convex PBW-type Lyndon basis of the algebra* U^+. □

Remark: (1) One can take a lexicographical ordering on the set of positive quantum root vectors provided that one puts $j := 2n - j' + 1$ for $1 \leq j' \leq n$. That is, the set

$$\left\{ \mathcal{E}_{i_p,j_p} \cdots \mathcal{E}_{i_2,j_2}\mathcal{E}_{i_1,j_1} \;\middle|\; (i_1, j_1) \leq (i_2, j_2) \leq \cdots \leq (i_p, j_p) \quad \textit{lexicographically} \right\},$$

where $1 \leq i_l \leq n$ and $1 \leq j_l \leq 2n$, is a basis of the algebra U^+.

(2) The set of elements $\mathcal{E}_{1,1}^{c_1}\mathcal{E}_{1,2}^{c_2}\mathcal{E}_{1,3}^{c_3} \cdots \mathcal{E}_{1,n}^{c_n}\mathcal{E}_{1,n'}^{c_{n+1}} \cdots \mathcal{E}_{1,2'}^{c_{2n-1}}\mathcal{E}_{2,2}^{c_{2n}}\mathcal{E}_{2,3}^{c_{2n+1}} \cdots \mathcal{E}_{2,3'}^{c_{4n-4}}$ $\mathcal{E}_{3,3}^{c_{4n-3}} \cdots \mathcal{E}_{n-1,n-1}^{c_{n^2-3}}\mathcal{E}_{n-1,n}^{c_{n^2-2}}\mathcal{E}_{n-1,n'}^{c_{n^2-1}}\mathcal{E}_{n,n}^{c_{n^2}}$ $\left((c_1, \cdots, c_{n^2}) \in N^{n^2}\right)$ forms a basis of U^+.

2.6. The following τ plays a key role in describing Lyndon basis (see [**HRZ**]).

Definition 2.5. Let τ be the $\mathbb{Q}$-algebra anti-automorphism of $U_{r,s}(\mathfrak{so}_{2n+1})$ such that $\tau(r) = s$, $\tau(s) = r$, $\tau(\langle\omega_i', \omega_j\rangle^{\pm 1}) = \langle\omega_j', \omega_i\rangle^{\mp 1}$, and

$$\tau(e_i) = f_i, \quad \tau(f_i) = e_i, \quad \tau(\omega_i) = \omega_i', \quad \tau(\omega_i') = \omega_i.$$

Then $\mathcal{B}' = \tau(\mathcal{B})$ with those induced defining relations from $\mathcal{B}$, and those cross relations in (B2)—(B4) are antisymmetric with respect to τ. □

Using τ to U^+, we can get those negative quantum root vectors in U^-. Define $\mathcal{F}_{i,i} = \tau(\mathcal{E}_{i,i}) = f_i$ for $1 \leq i \leq n$, and

$$\mathcal{F}_{i,j} = \tau(\mathcal{E}_{i,j}) = \mathcal{F}_{i+1,j}f_i - s^2 f_i\mathcal{F}_{i+1,j}, \qquad 1 \leq i < j \leq n, \tag{2.8}$$

$$\mathcal{F}_{i,n'} = \tau(\mathcal{E}_{i,n'}) = f_n\mathcal{F}_{i,n} - rs\mathcal{F}_{i,n}f_n, \qquad 1 \leq i \leq n-1, \tag{2.9}$$

$$\mathcal{F}_{i,j'} = \tau(\mathcal{E}_{i,j'}) = f_j\mathcal{F}_{i,j+1'} - r^{-2}\mathcal{F}_{i,j+1'}f_j, \quad 1 \leq i < j \leq n-1. \tag{2.10}$$

Corollary 2.6. *$\Big\{ \mathcal{F}_{1,1}^{c_1}\mathcal{F}_{1,2}^{c_2}\mathcal{F}_{1,3}^{c_3} \cdots \mathcal{F}_{1,n}^{c_n}\mathcal{F}_{1,n'}^{c_{n+1}} \cdots \mathcal{F}_{1,2'}^{c_{2n-1}} \cdot \mathcal{F}_{2,2}^{c_{2n}}\mathcal{F}_{2,3}^{c_{2n+1}} \cdots \mathcal{F}_{2,n}^{c_{3n-2}}$ $\mathcal{F}_{2,n'}^{c_{3n-1}} \cdots \mathcal{F}_{2,3'}^{c_{4n-4}} \cdots \cdot \mathcal{F}_{n-1,n-1}^{c_{n^2-3}}\mathcal{F}_{n-1,n}^{c_{n^2-2}}\mathcal{F}_{n-1,n'}^{c_{n^2-1}} \cdot \mathcal{F}_{n,n}^{c_{n^2}} \;\Big|\; (c_1, \cdots, c_{n^2}) \in \mathbb{Z}_+^{n^2} \Big\}$ forms a convex PBW-type basis of the algebra U^-.* □

3. Commutation relations and homogeneous central elements

3.1. Commutation relations in U^+. We will make more efforts to explicitly calculate all possible commutation relations, which are useful for determining central elements in subsection *3.2* and integrals in Section 7. We point out the fact that these commutation relations or say, (r,s)-identities hold in U^+ arises essentially from the (r,s)-Serre relations $(B5)$. First of all, we pay attention to the following

Question: Is the defining result of (r,s)-bracketings in *Remark* (ii) unique for those non-unique decomposition of $\gamma = \alpha' + \beta'$ such that $\alpha', \beta' \in \Phi^+$ and $\alpha' < \gamma < \beta'$? i.e., does it have $[\mathcal{E}_{\alpha'}, \mathcal{E}_{\beta'}]_{\langle\omega_{\alpha'}', \omega_{\beta'}\rangle^{-1}} = [\mathcal{E}_\alpha, \mathcal{E}_\beta]_{\langle\omega_\alpha', \omega_\beta\rangle^{-1}}$ for such a γ?

Lemma 3.1 (3) & (4) below have a positive answer to this question provided that we work under the structure constants matrix listed in subsection *2.1.* We will see that Lemma 3.1 (3) & (4) are not only the most basic ones in all commutation relations involved but also well-adopted to our calculating technique.

For simplicity, we write (r,s)-bracketings by $[\mathcal{E}_\alpha, \mathcal{E}_\beta]_\bullet = [\mathcal{E}_\alpha, \mathcal{E}_\beta]_{\langle\omega_\alpha', \omega_\beta\rangle^{-1}}$ briefly.

Lemma 3.1. *The following relations hold in U^+ :*

(1) $\mathcal{E}_{i,j}\mathcal{E}_{k,l} = \mathcal{E}_{k,l}\mathcal{E}_{i,j}$, $\qquad i \leq j,\ j+1 < k \leq l \leq n$;
(2) $\mathcal{E}_{i,j}\mathcal{E}_{k,l'} = \mathcal{E}_{k,l'}\mathcal{E}_{i,j}$, $\qquad i \leq j,\ j+1 < k < l \leq n$;
(3) $\mathcal{E}_{i,j} = \mathcal{E}_{i,l-1}\mathcal{E}_{l,j} - r^2\mathcal{E}_{l,j}\mathcal{E}_{i,l-1}$, $\qquad i < l \leq j \leq n$;
(4) $\mathcal{E}_{i,j'} = \mathcal{E}_{i,l-1}\mathcal{E}_{l,j'} - r^2\mathcal{E}_{l,j'}\mathcal{E}_{i,l-1}$, $\qquad i < l < j \leq n$.

Proof. (1) & (2) follow directly from the (r,s)-Serre relation $(B5)$.

(3): We fix l and j with $l \leq j$ and use induction on i. If $i = l-1$, this is just the defining formula (2.2) of $\mathcal{E}_{i,j}$. Assume that $i < l-1$, then we have

$$\begin{aligned}\mathcal{E}_{i,l-1}\mathcal{E}_{l,j} &= (e_i\mathcal{E}_{i+1,l-1} - r^2\mathcal{E}_{i+1,l-1}e_i)\mathcal{E}_{l,j}\\ &= e_i\mathcal{E}_{i+1,l-1}\mathcal{E}_{l,j} - r^2\mathcal{E}_{i+1,l-1}\mathcal{E}_{l,j}e_i\\ &= e_i(r^2\mathcal{E}_{l,j}\mathcal{E}_{i+1,l-1} + \mathcal{E}_{i+1,j}) - r^2(r^2\mathcal{E}_{l,j}\mathcal{E}_{i+1,l-1} + \mathcal{E}_{i+1,j})e_i\\ &= r^2\mathcal{E}_{l,j}e_i\mathcal{E}_{i+1,l-1} + e_i\mathcal{E}_{i+1,j} - r^4\mathcal{E}_{l,j}\mathcal{E}_{i+1,l-1}e_i - r^2\mathcal{E}_{i+1,j}e_i\\ &= r^2\mathcal{E}_{l,j}\mathcal{E}_{i,l-1} + \mathcal{E}_{i,j}\end{aligned}$$

by (1), the induction hypothesis and the defining formula (2.2).

(4): The following expression can be easily verified:

$$\mathcal{E}_{i,n'} = e_i\mathcal{E}_{i+1,n'} - r^2\mathcal{E}_{i+1,n'}e_i, \quad i \leq n-2. \tag{$*$}$$

Indeed, for $i \leq n-2$, we have $e_ie_n = e_ne_i$. By the defining relation (2.3), we get

$$\begin{aligned}e_i\mathcal{E}_{i+1,n'} &= e_i(\mathcal{E}_{i+1,n}e_n - rse_n\mathcal{E}_{i+1,n})\\ &= (\mathcal{E}_{i,n} + r^2\mathcal{E}_{i+1,n}e_i)e_n - rse_n(\mathcal{E}_{i,n} + r^2\mathcal{E}_{i+1,n}e_i)\\ &= \mathcal{E}_{i,n'} + r^2\mathcal{E}_{i+1,n'}e_i.\end{aligned}$$

We suppose initially that $j = n$ and $i = l-1$, this is just the relation in $(*)$.

Now assume (4) is true for $\mathcal{E}_{i+1,n'}$. This means $i+1 < l$. Then using (2), $e_i\mathcal{E}_{l,n'} = \mathcal{E}_{l,n'}e_i$. By the induction hypothesis and $(*)$, we obtain

$$\begin{aligned}\mathcal{E}_{i,l-1}\mathcal{E}_{l,n'} &= (e_i\mathcal{E}_{i+1,l-1} - r^2\mathcal{E}_{i+1,l-1}e_i)\mathcal{E}_{l,n'}\\ &= e_i(\mathcal{E}_{i+1,n'} + r^2\mathcal{E}_{l,n'}\mathcal{E}_{i+1,l-1}) - r^2\mathcal{E}_{i+1,l-1}\mathcal{E}_{l,n'}e_i\\ &= e_i\mathcal{E}_{i+1,n'} + r^2\mathcal{E}_{l,n'}e_i\mathcal{E}_{i+1,l-1} - r^2\mathcal{E}_{i+1,n'}e_i - r^4\mathcal{E}_{l,n'}\mathcal{E}_{i+1,l-1}e_i\\ &= \mathcal{E}_{i,n'} + r^2\mathcal{E}_{l,n'}\mathcal{E}_{i,l-1}.\end{aligned}$$

Finally, assume (4) is true for $\mathcal{E}_{i,j+1'}$ with $i < l$. Since $i \leq l-1 < l < j$, $e_j\mathcal{E}_{i,l-1} = \mathcal{E}_{i,l-1}e_j$ by (1). Using the defining formula (2.4) and the induction hypothesis, we get

$$\begin{aligned}\mathcal{E}_{i,j'} &= \mathcal{E}_{i,j+1'}e_j - s^{-2}e_j\mathcal{E}_{i,j+1'}\\ &= (\mathcal{E}_{i,l-1}\mathcal{E}_{l,j+1'} - r^2\mathcal{E}_{l,j+1'}\mathcal{E}_{i,l-1})e_j - s^{-2}e_j(\mathcal{E}_{i,l-1}\mathcal{E}_{l,j+1'} - r^2\mathcal{E}_{l,j+1'}\mathcal{E}_{i,l-1})\\ &= \mathcal{E}_{i,l-1}\mathcal{E}_{l,j+1'}e_j - r^2\mathcal{E}_{l,j+1'}e_j\mathcal{E}_{i,l-1} - s^{-2}\mathcal{E}_{i,l-1}e_j\mathcal{E}_{l,j+1'} + s^{-2}r^2e_j\mathcal{E}_{l,j+1'}\mathcal{E}_{i,l-1}\\ &= \mathcal{E}_{i,l-1}\mathcal{E}_{l,j'} - r^2\mathcal{E}_{l,j'}\mathcal{E}_{i,l-1}.\end{aligned}$$

Thus, the proof is complete. □

Lemma 3.2. *The following relations hold in* U^+ :

(1) $e_i\mathcal{E}_{i,j} = s^2\mathcal{E}_{i,j}e_i, \qquad 1 \leq i < j \leq n;$
(2) $e_i\mathcal{E}_{i,j'} = s^2\mathcal{E}_{i,j'}e_i, \qquad i+1 < j \leq n;$
(3) $\mathcal{E}_{i,j}e_j = s^2e_j\mathcal{E}_{i,j}, \qquad 1 \leq i < j < n;$
(4) $\mathcal{E}_{i,n'}e_n = s^2e_n\mathcal{E}_{i,n'}, \qquad 1 \leq i < n;$
(5) $\mathcal{E}_{i,j'}e_j = r^{-2}e_j\mathcal{E}_{i,j'}, \qquad 1 \leq i < j < n.$

Proof. (1) follows directly from the (r,s)-Serre relations $(B5)$, (2.5) and Lemma 3.1 (1) & (3) as one can decompose $\mathcal{E}_{i,j} = [\mathcal{E}_{i,i+1}, \mathcal{E}_{i+2,j}]_{r^2}$.

(2), (3) & (4) can be proved similarly using the (r,s)-Serre relations $(B5)$, (2.6) or (2.7), together with Lemma 3.1.

(5) will be proved in Lemma 3.6 (2). □

LEMMA 3.3. *The following relations hold in* U^+ :

(1) $e_l\mathcal{E}_{i,j}=\mathcal{E}_{i,j}e_l,\qquad i<l<j\le n;$
(2) $\mathcal{E}_{i,l}\mathcal{E}_{l,j}-(rs)^2\mathcal{E}_{l,j}\mathcal{E}_{i,l}=(s^2-r^2)e_l\mathcal{E}_{i,j},\qquad i<l<j\le n;$
(3) $\mathcal{E}_{i,j}\mathcal{E}_{k,l}=\mathcal{E}_{k,l}\mathcal{E}_{i,j},\qquad i<k\le l<j\le n;$
(4) $\mathcal{E}_{i,j}\mathcal{E}_{l,j}=s^2\mathcal{E}_{l,j}\mathcal{E}_{i,j},\qquad i<l\le j<n;$
(5) $\mathcal{E}_{i,l}\mathcal{E}_{i,j}=s^2\mathcal{E}_{i,j}\mathcal{E}_{i,l},\qquad i\le l<j\le n;$
(6) $e_l\mathcal{E}_{i,j'}=\mathcal{E}_{i,j'}e_l,\qquad i<l,l+1<j\le n;$
(7) $\mathcal{E}_{i,l}\mathcal{E}_{l,j'}-(rs)^2\mathcal{E}_{l,j'}\mathcal{E}_{i,l}=(s^2-r^2)e_l\mathcal{E}_{i,j'},\qquad i<l,l+1<j\le n;$
(8) $\mathcal{E}_{i,j'}\mathcal{E}_{k,l}=\mathcal{E}_{k,l}\mathcal{E}_{i,j'},\qquad i<k\le l,l+1<j\le n;$
(9) $\mathcal{E}_{i,l}\mathcal{E}_{i,j'}=s^2\mathcal{E}_{i,j'}\mathcal{E}_{i,l},\qquad i\le l,l+1<j\le n.$

PROOF. (1) & (2): Using Lemma 3.1 (3) to $\mathcal{E}_{l,j}$ and Lemma 3.2 (3), we get

$$\begin{aligned}
&\mathcal{E}_{i,l}\mathcal{E}_{l,j}-(rs)^2\mathcal{E}_{l,j}\mathcal{E}_{i,l}\\
&\quad=\mathcal{E}_{i,l}(e_l\mathcal{E}_{l+1,j}-r^2\mathcal{E}_{l+1,j}e_l)-(rs)^2(e_l\mathcal{E}_{l+1,j}-r^2\mathcal{E}_{l+1,j}e_l)\mathcal{E}_{i,l}\\
&\quad=s^2e_l\mathcal{E}_{i,l}\mathcal{E}_{l+1,j}-r^2\mathcal{E}_{i,l}\mathcal{E}_{l+1,j}e_l-(rs)^2e_l\mathcal{E}_{l+1,j}\mathcal{E}_{i,l}+r^4\mathcal{E}_{l+1,j}\mathcal{E}_{i,l}e_l\\
&\quad=s^2e_l\mathcal{E}_{i,j}-r^2\mathcal{E}_{i,j}e_l.
\end{aligned}$$

Again, using Lemma 3.1 (3) to $\mathcal{E}_{i,l}$ and Lemma 3.2 (1), we have

$$\begin{aligned}
&\mathcal{E}_{i,l}\mathcal{E}_{l,j}-(rs)^2\mathcal{E}_{l,j}\mathcal{E}_{i,l}\\
&\quad=(\mathcal{E}_{i,l-1}e_l-r^2e_l\mathcal{E}_{i,l-1})\mathcal{E}_{l,j}-(rs)^2\mathcal{E}_{l,j}(\mathcal{E}_{i,l-1}e_l-r^2e_l\mathcal{E}_{i,l-1})\\
&\quad=s^2\mathcal{E}_{i,l-1}\mathcal{E}_{l,j}e_l-r^2e_l\mathcal{E}_{i,l-1}\mathcal{E}_{l,j}-(rs)^2\mathcal{E}_{l,j}\mathcal{E}_{i,l-1}e_l+r^4e_l\mathcal{E}_{l,j}\mathcal{E}_{i,l-1}\\
&\quad=s^2\mathcal{E}_{i,j}e_l-r^2e_l\mathcal{E}_{i,j}.
\end{aligned}$$

Combining the above results, we have $(r^2+s^2)e_l\mathcal{E}_{i,j}=(r^2+s^2)\mathcal{E}_{i,j}e_l$. By assumption $r^2+s^2\neq 0$, we obtain (1), hence (2).

(3) is a direct consequence of (1).

(4) follows from (3) and Lemma 3.2 (3) since $\mathcal{E}_{l,j}=[\mathcal{E}_{l,j-1},e_j]_{r^2}$ by Lemma 3.1 (3).

(5) can be proved similarly to (4), by (3), Lemma 3.1 (3) and Lemma 3.2 (1).

(6), (7): can be proved similarly to (1) & (2), respectively.

(8) follows from (6).

(9) follows from (8), together with Lemma 3.2 (2). □

LEMMA 3.4. *The following relations hold in* U^+ :

(1) $\mathcal{E}_{i,n}\mathcal{E}_{j,n}-(rs)\mathcal{E}_{j,n}\mathcal{E}_{i,n}=\mathcal{E}_{j,n-1}\mathcal{E}_{i,n'}-r^2\mathcal{E}_{i,n'}\mathcal{E}_{j,n-1},\qquad i<j<n;$
(2) $\mathcal{E}_{j,k}\mathcal{E}_{i,k+1'}-r^2\mathcal{E}_{i,k+1'}\mathcal{E}_{j,k}=\mathcal{E}_{i,k+2'}\mathcal{E}_{j,k+1}-s^{-2}\mathcal{E}_{j,k+1}\mathcal{E}_{i,k+2'},\qquad i<j<k<n-1;$
(3) $\mathcal{E}_{n-1,n}\mathcal{E}_{n-1,n'}=s^2\mathcal{E}_{n-1,n'}\mathcal{E}_{n-1,n};$
(4) $\mathcal{E}_{i,j'}e_n=(rs)^2e_n\mathcal{E}_{i,j'},\qquad i<j<n;$
(5) $\mathcal{E}_{i,n'}\mathcal{E}_{n-1,n}=\mathcal{E}_{n-1,n}\mathcal{E}_{i,n'},\qquad i<n-1;$
(6) $\mathcal{E}_{i,n'}\mathcal{E}_{n-1,n'}=s^2\mathcal{E}_{n-1,n'}\mathcal{E}_{i,n'},\qquad i<n-1;$
(7) $\mathcal{E}_{i,n-1'}\mathcal{E}_{n-1,n}=s^2\mathcal{E}_{n-1,n}\mathcal{E}_{i,n-1'},\qquad i<n-1;$
(8) $\mathcal{E}_{i,n-1'}\mathcal{E}_{n-1,n'}=(rs^2)^2\mathcal{E}_{n-1,n'}\mathcal{E}_{i,n-1'},\qquad i<n-1.$

PROOF. (1): By Lemma 3.3 (3), we have $\mathcal{E}_{i,n}\mathcal{E}_{j,n-1} = \mathcal{E}_{j,n-1}\mathcal{E}_{i,n}$. So by Lemma 3.1 (3), we obtain

$$\begin{aligned}
\mathcal{E}_{i,n}\mathcal{E}_{j,n} &= \mathcal{E}_{i,n}(\mathcal{E}_{j,n-1}e_n - r^2 e_n\mathcal{E}_{j,n-1}) \\
&= \mathcal{E}_{j,n-1}\mathcal{E}_{i,n}e_n - r^2\mathcal{E}_{i,n}e_n\mathcal{E}_{j,n-1} \\
&= \mathcal{E}_{j,n-1}\mathcal{E}_{i,n'} + rs\mathcal{E}_{j,n-1}e_n\mathcal{E}_{i,n} - r^2\mathcal{E}_{i,n'}\mathcal{E}_{j,n-1} - r^3 s e_n\mathcal{E}_{i,n}\mathcal{E}_{j,n-1} \\
&= \mathcal{E}_{j,n-1}\mathcal{E}_{i,n'} + rs\mathcal{E}_{j,n}\mathcal{E}_{i,n} - r^2\mathcal{E}_{i,n'}\mathcal{E}_{j,n-1}.
\end{aligned}$$

(2): By Lemma 3.3 (8), we have $\mathcal{E}_{j,k}\mathcal{E}_{i,k+2'} = \mathcal{E}_{i,k+2'}\mathcal{E}_{j,k}$. Using $\mathcal{E}_{j,k+1} = \mathcal{E}_{j,k}e_{k+1} - r^2 e_{k+1}\mathcal{E}_{j,k}$, we obtain

$$\begin{aligned}
\mathcal{E}_{j,k}\mathcal{E}_{i,k+1'} &= \mathcal{E}_{j,k}\Big(\mathcal{E}_{i,k+2'}e_{k+1} - s^{-2}e_{k+1}\mathcal{E}_{i,k+2'}\Big) \\
&= \mathcal{E}_{i,k+2'}\mathcal{E}_{j,k}e_{k+1} - s^{-2}\mathcal{E}_{j,k}e_{k+1}\mathcal{E}_{i,k+2'}) \\
&= \mathcal{E}_{i,k+2'}\mathcal{E}_{j,k+1} + r^2\mathcal{E}_{i,k+2'}e_{k+1}\mathcal{E}_{j,k} \\
&\quad - s^{-2}\mathcal{E}_{j,k+1}\mathcal{E}_{i,k+2'} - r^2 s^{-2}e_{k+1}\mathcal{E}_{i,k+2'}\mathcal{E}_{j,k} \\
&= r^2\mathcal{E}_{i,k+1'}\mathcal{E}_{j,k} + \mathcal{E}_{i,k+2'}\mathcal{E}_{j,k+1} - s^{-2}\mathcal{E}_{j,k+1}\mathcal{E}_{i,k+2'}.
\end{aligned}$$

(3) can be proved by the same argument used in checking the commutation relation (6) of Lemma 3.6 in [**BGH1**] (where the identity $\mathcal{E}_{1'2}\mathcal{E}_{12} = r^2\mathcal{E}_{12}\mathcal{E}_{1'2}$ was actually verified for type C_2 case by only using (r,s)-Serre relations).

(4): It suffices to check the case when $j = n-1$, i.e., $\mathcal{E}_{i,n-1'}e_n = (rs)^2 e_n\mathcal{E}_{i,n-1'}$, since for any $j \le n-2$, $e_n e_j = e_j e_n$ and $\mathcal{E}_{i,j'} = [\cdots[\,\mathcal{E}_{i,n-1'}, e_{n-2}]_{s^{-2}}, \cdots, e_j\,]_{s^{-2}}$, then $\mathcal{E}_{i,j'}e_n = (rs)^2 e_n\mathcal{E}_{i,j'}$.

Observe that (4) in the case when $j = n-1$ is equivalent to (5):

$$\begin{aligned}
&\mathcal{E}_{i,n-1'}e_n - (rs)^2 e_n\mathcal{E}_{i,n-1'} \\
&\quad = \big(\mathcal{E}_{i,n'}e_{n-1} - s^{-2}e_{n-1}\mathcal{E}_{i,n'}\big)e_n - (rs)^2 e_n\big(\mathcal{E}_{i,n'}e_{n-1} - s^{-2}e_{n-1}\mathcal{E}_{i,n'}\big) \\
&\quad = \mathcal{E}_{i,n'}e_{n-1}e_n - e_{n-1}e_n\mathcal{E}_{i,n'} - r^2\mathcal{E}_{i,n'}e_n e_{n-1} + r^2 e_n e_{n-1}\mathcal{E}_{i,n'} \\
&\quad = \mathcal{E}_{i,n'}\mathcal{E}_{n-1,n} - \mathcal{E}_{n-1,n}\mathcal{E}_{i,n'}.
\end{aligned}$$

Since for $i < n-2$, we have the decompositions $\mathcal{E}_{i,n-1'} = [\,\mathcal{E}_{i,n-3}, \mathcal{E}_{n-2,n-1'}]_{r^2}$ and $\mathcal{E}_{i,n'} = [\,\mathcal{E}_{i,n-3}, \mathcal{E}_{n-2,n'}]_{r^2}$, by Lemma 3.1 (4). Thus, this reduces to verify $\mathcal{E}_{n-2,n-1'}e_n = (rs)^2 e_n\mathcal{E}_{n-2,n-1'}$ or equivalently, $\mathcal{E}_{n-2,n'}\mathcal{E}_{n-1,n} = \mathcal{E}_{n-1,n}\mathcal{E}_{n-2,n'}$. While the latter is equivalent to (3) under assumption $r^3 \ne s^3$.

Indeed, using (2.3) and (1), we at first have

$$\begin{aligned}
&\mathcal{E}_{n-2,n}\mathcal{E}_{n-1,n'} - (rs)^2\mathcal{E}_{n-1,n'}\mathcal{E}_{n-2,n} \\
&\quad = \big(\mathcal{E}_{n-2,n}\mathcal{E}_{n-1,n}\big)e_n - rs\big(\mathcal{E}_{n-2,n}e_n\big)\mathcal{E}_{n-1,n} - (rs)^2\mathcal{E}_{n-1,n'}\mathcal{E}_{n-2,n} \qquad \text{(by (1), (2.3))} \\
&\quad = \Big(rs\mathcal{E}_{n-1,n}\mathcal{E}_{n-2,n} + e_{n-1}\mathcal{E}_{n-2,n'} - r^2\mathcal{E}_{n-2,n'}e_{n-1}\Big)e_n \\
&\qquad - rs\mathcal{E}_{n-2,n'}\mathcal{E}_{n-1,n} - (rs)^2 e_n\big(\mathcal{E}_{n-2,n}\mathcal{E}_{n-1,n}\big) - (rs)^2\mathcal{E}_{n-1,n'}\mathcal{E}_{n-2,n}
\end{aligned}$$

$$\begin{aligned}
&= rs\mathcal{E}_{n-1,n}\big(\mathcal{E}_{n-2,n'}+\underline{rse_n\mathcal{E}_{n-2,n}}\big)+s^2e_{n-1}e_n\mathcal{E}_{n-2,n'}-r^2\mathcal{E}_{n-2,n'}e_{n-1}e_n\\
&\quad -rs\mathcal{E}_{n-2,n'}\mathcal{E}_{n-1,n}-(rs)^2e_n\Big(\underline{rs\mathcal{E}_{n-1,n}\mathcal{E}_{n-2,n}}+e_{n-1}\mathcal{E}_{n-2,n'}-r^2\mathcal{E}_{n-2,n'}e_{n-1}\Big)\\
&\quad -\underline{(rs)^2\mathcal{E}_{n-1,n'}\mathcal{E}_{n-2,n}} \qquad\qquad \text{(applying (2.3) to the terms underlined)}\\
&= rs\mathcal{E}_{n-1,n}\mathcal{E}_{n-2,n'}+s^2e_{n-1}e_n\mathcal{E}_{n-2,n'}-r^2\mathcal{E}_{n-2,n'}e_{n-1}e_n\\
&\quad -rs\mathcal{E}_{n-2,n'}\mathcal{E}_{n-1,n}-(rs)^2e_ne_{n-1}\mathcal{E}_{n-2,n'}+r^4\mathcal{E}_{n-2,n'}e_ne_{n-1}\\
&= (rs+s^2)\mathcal{E}_{n-1,n}\mathcal{E}_{n-2,n'}-(r^2+rs)\mathcal{E}_{n-2,n'}\mathcal{E}_{n-1,n}.
\end{aligned}$$

Next, using Lemma 3.1 (3), together with (3): $\mathcal{E}_{n-1,n}\mathcal{E}_{n-1,n'} = s^2\mathcal{E}_{n-1,n'}\mathcal{E}_{n-1,n}$ and Lemma 3.1 (4), we have another expansion

$$\begin{aligned}
&\mathcal{E}_{n-2,n}\mathcal{E}_{n-1,n'}-(rs)^2\mathcal{E}_{n-1,n'}\mathcal{E}_{n-2,n}\\
&\quad = e_{n-2}\mathcal{E}_{n-1,n}\mathcal{E}_{n-1,n'} - r^2\mathcal{E}_{n-1,n}e_{n-2}\mathcal{E}_{n-1,n'}\\
&\qquad - (rs)^2\mathcal{E}_{n-1,n'}e_{n-2}\mathcal{E}_{n-1,n} + (r^2s)^2\mathcal{E}_{n-1,n'}\mathcal{E}_{n-1,n}e_{n-2}\\
&\quad = s^2e_{n-2}\mathcal{E}_{n-1,n'}\mathcal{E}_{n-1,n} - r^2\mathcal{E}_{n-1,n}e_{n-2}\mathcal{E}_{n-1,n'}\\
&\qquad - (rs)^2\mathcal{E}_{n-1,n'}e_{n-2}\mathcal{E}_{n-1,n} + r^4\mathcal{E}_{n-1,n}\mathcal{E}_{n-1,n'}e_{n-2}\\
&\quad = s^2\mathcal{E}_{n-2,n'}\mathcal{E}_{n-1,n} - r^2\mathcal{E}_{n-1,n}\mathcal{E}_{n-2,n'}.
\end{aligned}$$

Combining the above two expansions, we obtain

$$\mathcal{E}_{n-2,n'}\mathcal{E}_{n-1,n} = \mathcal{E}_{n-1,n}\mathcal{E}_{n-2,n'},$$

under assumption $r^2 + rs + s^2 \neq 0$.

(6) & (7) follow from (5), owing to (2.5) & (2.7).

(8) follows from (7) & (4). □

Lemma 3.5. *The following relations hold in* U^+ :

$$\begin{array}{lll}
(1) & \mathcal{E}_{i,n'}\mathcal{E}_{j,n} = \mathcal{E}_{j,n}\mathcal{E}_{i,n'}, & 1 \le i < j < n;\\
(2) & \mathcal{E}_{i,n}\mathcal{E}_{j,n'} - (rs)^2\mathcal{E}_{j,n'}\mathcal{E}_{i,n} = (s^2-r^2)\mathcal{E}_{i,n'}\mathcal{E}_{j,n}, & 1 \le i < j < n;\\
(3) & \mathcal{E}_{i,n}\mathcal{E}_{i,j'} = s^2\mathcal{E}_{i,j'}\mathcal{E}_{i,n}, & 1 \le i < j \le n;\\
(4) & \mathcal{E}_{i,n'}\mathcal{E}_{j,n'} = s^2\mathcal{E}_{j,n'}\mathcal{E}_{i,n'}, & 1 \le i < j < n;\\
(5) & \mathcal{E}_{i,l'}\mathcal{E}_{j,n} = \mathcal{E}_{j,n}\mathcal{E}_{i,l'}, & i < j < l \le n;\\
(6) & \mathcal{E}_{i,l'}\mathcal{E}_{j,n'} = (rs)^2\mathcal{E}_{j,n'}\mathcal{E}_{i,l'}, & i < j < l < n.
\end{array}$$

Proof. (1): Note that $\mathcal{E}_{i,n'}\mathcal{E}_{n-1,n} = \mathcal{E}_{n-1,n}\mathcal{E}_{i,n'}$ by Lemma 3.4 (5). So for $j \le n-2$, $\mathcal{E}_{i,n'}\mathcal{E}_{j,n-2} = \mathcal{E}_{j,n-2}\mathcal{E}_{i,n'}$ since $\mathcal{E}_{i,n'} = [\mathcal{E}_{i,n}, e_n]_{rs}$ and both $\mathcal{E}_{i,n}$ and e_n commute with $\mathcal{E}_{j,n-2}$ (by Lemma 3.3 (3)). Hence, Lemma 3.1 (3) gives us a decomposition $\mathcal{E}_{j,n} = [\mathcal{E}_{j,n-2}, \mathcal{E}_{n-1,n}]_{r^2}$, from which we obtain the required result.

(2): By (2.3), and using Lemma 3.2 (4), Lemma 3.4 (1), we have

$$\begin{aligned}
&\mathcal{E}_{i,n}\mathcal{E}_{j,n'}-(rs)^2\mathcal{E}_{j,n'}\mathcal{E}_{i,n}\\
&\quad = (\mathcal{E}_{i,n}\mathcal{E}_{j,n})e_n-rs(\mathcal{E}_{i,n}e_n)\mathcal{E}_{j,n}-(rs)^2\mathcal{E}_{j,n}(e_n\mathcal{E}_{i,n}) + (rs)^3e_n\mathcal{E}_{j,n}\mathcal{E}_{i,n}\\
&\quad = rs\mathcal{E}_{j,n}\mathcal{E}_{i,n}e_n+\mathcal{E}_{j,n-1}\mathcal{E}_{i,n'}e_n-r^2\mathcal{E}_{i,n'}\mathcal{E}_{j,n-1}e_n\\
&\qquad -rs(\mathcal{E}_{i,n'}+rse_n\mathcal{E}_{i,n})\mathcal{E}_{j,n}-(rs)^2\mathcal{E}_{j,n}(e_n\mathcal{E}_{i,n})+(rs)^3e_n\mathcal{E}_{j,n}\mathcal{E}_{i,n}
\end{aligned}$$

$$
\begin{aligned}
&= rs\mathcal{E}_{j,n}\mathcal{E}_{i,n'} - rs\mathcal{E}_{i,n'}\mathcal{E}_{j,n} + s^2\mathcal{E}_{j,n-1}e_n\mathcal{E}_{i,n'} - r^2\mathcal{E}_{i,n'}\mathcal{E}_{j,n-1}e_n\\
&\qquad -(rs)^2e_n(\mathcal{E}_{i,n}\mathcal{E}_{j,n} - rs\mathcal{E}_{j,n}\mathcal{E}_{i,n})\\
&= rs\mathcal{E}_{j,n}\mathcal{E}_{i,n'} - rs\mathcal{E}_{i,n'}\mathcal{E}_{j,n} + s^2\mathcal{E}_{j,n-1}e_n\mathcal{E}_{i,n'} - r^2\mathcal{E}_{i,n'}\mathcal{E}_{j,n-1}e_n\\
&\qquad -(rs)^2e_n(\mathcal{E}_{j,n-1}\mathcal{E}_{i,n'} - r^2\mathcal{E}_{i,n'}\mathcal{E}_{j,n-1})\\
&= rs[\mathcal{E}_{j,n}, \mathcal{E}_{i,n'}] + s^2\mathcal{E}_{j,n}\mathcal{E}_{i,n'} - r^2\mathcal{E}_{i,n'}\mathcal{E}_{j,n}
\end{aligned}
$$

which gives the required result using (1).

(3): First of all, we claim that $\mathcal{E}_{i,n}\mathcal{E}_{i,n'} = s^2\mathcal{E}_{i,n'}\mathcal{E}_{i,n}$. When $i = n-1$, this is Lemma 3.4 (3). For $i < n-1$, it follows from (1) since $\mathcal{E}_{i+1,n}\mathcal{E}_{i,n'} = \mathcal{E}_{i,n'}\mathcal{E}_{i+1,n}$ and $e_i\mathcal{E}_{i,n'} = s^2\mathcal{E}_{i,n'}e_i$, as well as $\mathcal{E}_{i,n} = [\,e_i, \mathcal{E}_{i+1,n}]_{r^2}$.

Next, noting $\mathcal{E}_{i,j'} = [\cdots[\,\mathcal{E}_{i,n'}, e_{n-1}]_{s^{-2}}, \cdots, e_j]_{s^{-2}}$ and $\mathcal{E}_{i,n}e_k = e_k\mathcal{E}_{i,n}$ for $i < j \le k < n$, together with the above assertion, we obtain the required result.

(4) follows from (1), Lemma 3.2 (4) as well as $\mathcal{E}_{j,n'} = [\,\mathcal{E}_{j,n}, e_n]_{rs}$.

(5) follows from (1) and $\mathcal{E}_{i,l'} = [\cdots[\,\mathcal{E}_{i,n'}, e_{n-1}]_{s^{-2}}, \cdots, e_l]_{s^{-2}}$, as well as $e_k\mathcal{E}_{j,n} = \mathcal{E}_{j,n}e_k$ for any k with $j < l \le k < n$.

(6) follows from (5) & Lemma 3.4 (4). □

Lemma 3.6. *The following relations hold in* U^+ :

(1) $[\,\mathcal{E}_{i,j'}, e_{i+1}]_\bullet = 0,\qquad 1 \le i < j \le n,\ i \ne j-2;$
(2) $\mathcal{E}_{i,j'}e_j = r^{-2}e_j\mathcal{E}_{i,j'},\qquad 1 \le i < j < n;$
(3) $[\,\mathcal{E}_{i,i+1'}, e_{i+2}]_\bullet = 0,\qquad 1 \le i < n-1;$
(4) $\mathcal{E}_{i,j'}e_l = e_l\mathcal{E}_{i,j'},\qquad i < j < l < n;$
(5) $\mathcal{E}_{i,l'}\mathcal{E}_{i,j'} = r^{-2}\mathcal{E}_{i,j'}\mathcal{E}_{i,l'},\qquad i < j < l \le n;$
(6) $\mathcal{E}_{i,l'}\mathcal{E}_{j,k'} = (rs)^2\mathcal{E}_{j,k'}\mathcal{E}_{i,l'},\qquad i < j < l < k \le n;$
(7) $\mathcal{E}_{i,l'}\mathcal{E}_{j,l'} = s^2\mathcal{E}_{j,l'}\mathcal{E}_{i,l'},\qquad i < j < l < n;$
(8) $\mathcal{E}_{i,n-1}\mathcal{E}_{i,n'} - (rs)^2\mathcal{E}_{i,n'}\mathcal{E}_{i,n-1} = s(s-r)\mathcal{E}_{i,n}^2,\qquad 1 \le i < n.$

Proof. See the appendix. □

Lemma 3.7. *The following relations hold in* U^+ :

(1) $\mathcal{E}_{i,k}\mathcal{E}_{i,k+1'} - (rs)^2\mathcal{E}_{i,k+1'}\mathcal{E}_{i,k} = s^2\mathcal{E}_{i,k+2'}\mathcal{E}_{i,k+1} - s^{-2}\mathcal{E}_{i,k+1}\mathcal{E}_{i,k+2'},\quad k < n-1;$
(2) $\mathcal{E}_{j,j+1}\mathcal{E}_{j,j+1'} = (rs^2)^2\mathcal{E}_{j,j+1'}\mathcal{E}_{j,j+1},\qquad j < n-1;$
(3) $\mathcal{E}_{j-1,j'}\mathcal{E}_{j,j+1'} = (rs^2)^2\mathcal{E}_{j,j+1'}\mathcal{E}_{j-1,j'},\qquad j < n-1;$
(4) $\mathcal{E}_{i,j'}\mathcal{E}_{j,j+1'} = (rs^2)^2\mathcal{E}_{j,j+1'}\mathcal{E}_{i,j'},\qquad i < j < n-1;$
(5) $\mathcal{E}_{j,k}\mathcal{E}_{j,j+1'} = (rs^2)^2\mathcal{E}_{j,j+1'}\mathcal{E}_{j,k},\qquad j < k < n;$
(6) $\mathcal{E}_{j-1,j}\mathcal{E}_{i,j'} = (rs)^2\mathcal{E}_{i,j'}\mathcal{E}_{j-1,j},\qquad i+1 < j < n;$
(7) $\mathcal{E}_{l,k}\mathcal{E}_{i,j'} = (rs)^2\mathcal{E}_{i,j'}\mathcal{E}_{l,k},\qquad i < l < j \le k < n;$
(8) $\mathcal{E}_{i,k}\mathcal{E}_{i,j'} = (rs^2)^2\mathcal{E}_{i,j'}\mathcal{E}_{i,k},\qquad i < j \le k < n.$

Proof. (1): The proof is similar to that of Lemma 3.4 (2) and noting that $\mathcal{E}_{i,k}\mathcal{E}_{i,k+2'} = s^2\mathcal{E}_{i,k+2'}\mathcal{E}_{i,k}$.

(2): Using (1) and Lemma 3.2 (5), we obtain

$$
\begin{aligned}
\mathcal{E}_{j,j+1}\mathcal{E}_{j,j+1'} &= (e_je_{j+1} - r^2e_{j+1}e_j)\mathcal{E}_{j,j+1'}\\
&= r^2e_j\mathcal{E}_{j,j+1'}e_{j+1} - r^2e_{j+1}e_j\mathcal{E}_{j,j+1'}\\
&= r^2\Big((rs)^2\mathcal{E}_{j,j+1'}e_j + s^2\mathcal{E}_{j,j+2'}\mathcal{E}_{j,j+1} - s^{-2}\mathcal{E}_{j,j+1}\mathcal{E}_{j,j+2'}\Big)e_{j+1}\\
&\quad - r^2e_{j+1}\Big((rs)^2\mathcal{E}_{j,j+1'}e_j + s^2\mathcal{E}_{j,j+2'}\mathcal{E}_{j,j+1} - s^{-2}\mathcal{E}_{j,j+1}\mathcal{E}_{j,j+2'}\Big)\\
&= (r^2s)^2\mathcal{E}_{j,j+1'}\mathcal{E}_{j,j+1} + (rs^2)^2\mathcal{E}_{j,j+1'}\mathcal{E}_{j,j+1} - (rs^{-1})^2\mathcal{E}_{j,j+1}\mathcal{E}_{j,j+1'}.
\end{aligned}
$$

So we have

$$(1+r^2s^{-2})\mathcal{E}_{j,j+1}\mathcal{E}_{j,j+1'}=(rs^2)^2(1+r^2s^{-2})\mathcal{E}_{j,j+1'}\mathcal{E}_{j,j+1},$$

since $1+r^2s^{-2}\neq 0$, we get

$$\mathcal{E}_{j,j+1}\mathcal{E}_{j,j+1'}=(rs^2)^2\mathcal{E}_{j,j+1'}\mathcal{E}_{j,j+1}.$$

(3): By Lemma 3.6 (2), (4) & Lemma 3.4 (4), we have

$$\begin{aligned}\mathcal{E}_{j-1,j'}\mathcal{E}_{j,j+2'}&=\mathcal{E}_{j-1,j'}\big(e_j\mathcal{E}_{j+1,j+2'}-r^2\mathcal{E}_{j+1,j+2'}e_j\big)\\&=(rs^2)^2\big(e_j\mathcal{E}_{j+1,j+2'}-r^2\mathcal{E}_{j+1,j+2'}e_j\big)\mathcal{E}_{j-1,j'}\\&=(rs^2)^2\mathcal{E}_{j,j+2'}\mathcal{E}_{j-1,j'}.\end{aligned}$$

Again, by Lemma 3.6 (4), we arrive at

$$\begin{aligned}\mathcal{E}_{j-1,j'}\mathcal{E}_{j,j+1'}&=\mathcal{E}_{j-1,j'}\big(\mathcal{E}_{j,j+2'}e_{j+1}-s^{-2}e_{j+1}\mathcal{E}_{j,j+2'}\big)\\&=(rs^2)^2\big(\mathcal{E}_{j,j+2'}e_{j+1}-s^{-2}e_{j+1}\mathcal{E}_{j,j+2'}\big)\mathcal{E}_{j-1,j'}\\&=(rs^2)^2\mathcal{E}_{j,j+1'}\mathcal{E}_{j-1,j'}.\end{aligned}$$

(4) follows from (3), together with $e_l\mathcal{E}_{j,j+1'}=\mathcal{E}_{j,j+1'}e_l$ for $i\le l<j-1$.

(5) follows from (2), together with Lemma 3.6 (4).

(6) An observation when $i=j-2$ comes from the proof of Lemma 3.6 (3), for $j<n$. For $i<j-2$, it follows from Lemma 3.1 (1).

(7) follows from (6), together with Lemmas 3.3 (6) & 3.6 (4).

(8) follows from (7), together with Lemma 3.2 (2) for $i+1<j$, but the case for $i+1=j$ follows from (2) as well as Lemma 3.6 (4). □

3.2. Central elements. Restrict two parameters r and s to be roots of 1: r is a primitive dth root of unity, s is a primitive d'th root of unity and ℓ is the least common multiple of d and d'. From now on, we assume that $\mathbb{K}$ contains a primitive ℓth root of unity.

The following Lemma is useful to derive some commutator relations.

Lemma 3.8. *Let x,y,z be elements of a $\mathbb{K}$-algebra such that $yx=\alpha xy+z$ for some $\alpha\in\mathbb{K}$ and m a natural number. Then the following assertions hold*

(1) *If $zx=\beta xz$ for some $\beta(\neq\alpha)\in\mathbb{K}$, then $yx^m=\alpha^m x^m y+\frac{\alpha^m-\beta^m}{\alpha-\beta}x^{m-1}z$;*

(2) *If $yz=\beta zy$ for some $\beta(\neq\alpha)\in\mathbb{K}$, then $y^m x=\alpha^m xy^m+\frac{\alpha^m-\beta^m}{\alpha-\beta}zy^{m-1}$.*

Based on Lemma 3.8 and by induction, one can easily check the following Lemmas, where the (r,s)-integers, factorials and binomial coefficients are defined for positive integers c and d by

$$[c]_i:=\frac{r_i^c-s_i^c}{r_i-s_i},\quad [c]_i!:=[c]_i[c-1]_i\cdots[2]_i[1]_i,\quad \begin{bmatrix}c\\d\end{bmatrix}_i:=\frac{[c]_i!}{[d]_i![c-d]_i!}.$$

Write in brief $[m]=[m]_i$ for $i<n$. By convention $[0]_i=0$ and $[0]_i!=1$. Denote $\alpha_m=\frac{(r^m-s^m)(r^{m-1}-s^{m-1})}{(r-s)(r^2-s^2)}$, $\beta_m=\frac{r^m-s^m}{r-s}$, then $\alpha_{m+1}=s^2\alpha_m+r^{m-1}\beta_m$ and $\beta_{m+1}=s\beta_m+r^m$.

For $\mathcal{E}_{i,j}^m$ $(i\le j\le n)$ and $\mathcal{E}_{i,n'}^m$ $(i<n)$ with $m\in\mathbb{Z}_+$, we have

Lemma 3.9. *For $m\in\mathbb{Z}_+$ and $i\le j<n$, the following equalities hold*

(1) $e_{i-1}\mathcal{E}_{i,j}^m=r^{2m}\mathcal{E}_{i,j}^m e_{i-1}+[m]\ \mathcal{E}_{i,j}^{m-1}\mathcal{E}_{i-1,j}$;

(2) $\mathcal{E}_{i,j}^m e_{j+1}=r^{2m}e_{j+1}\mathcal{E}_{i,j}^m+[m]\ \mathcal{E}_{i,j+1}\mathcal{E}_{i,j}^{m-1}$;

(3) $e_{n-1}e_n^m = \alpha_m e_n^{m-2}\mathcal{E}_{n-1,n'} + r^{m-1}\beta_m e_n^{m-1}\mathcal{E}_{n-1,n} + r^{2m}e_n^m e_{n-1};$

(4) $\mathcal{E}_{i,n}^m e_n = (rs)^m e_n\mathcal{E}_{i,n}^m + s^{m-1}\beta_m\ \mathcal{E}_{i,n'}\mathcal{E}_{i,n}^{m-1};$

(5) $$e_{i-1}\mathcal{E}_{i,n}^m = r^{m-1}\beta_m\ \mathcal{E}_{i,n}^{m-1}\mathcal{E}_{i-1,n} + \alpha_m\ \mathcal{E}_{i,n}^{m-2}\mathcal{E}_{i,n-1}\mathcal{E}_{i-1,n'}$$
$$-r^2\alpha_m\ \mathcal{E}_{i,n}^{m-2}\mathcal{E}_{i-1,n'}\mathcal{E}_{i,n-1} + r^{2m}\mathcal{E}_{i,n}^m e_{i-1};$$

(6) $e_{n-1}\mathcal{E}_{n-1,n'}^m = s^{2m-1}(s-r)[m]\ \mathcal{E}_{n-1,n'}^{m-1}\mathcal{E}_{n-1,n}^2 + (rs)^{2m}\mathcal{E}_{n-1,n'}^m e_{n-1};$

(7) $\mathcal{E}_{i,n'}^m e_{n-1} = s^{2(1-m)}[m]\ \mathcal{E}_{i,n'}^{m-1}\mathcal{E}_{i,n-1'} + s^{-2m}e_{n-1}\mathcal{E}_{i,n'}^m, \qquad i < n-1;$

(8) $e_{i-1}\mathcal{E}_{i,n'}^m = [m]\ \mathcal{E}_{i,n'}^{m-1}\mathcal{E}_{i-1,n'} + r^{2m}\mathcal{E}_{i,n'}^m e_{i-1}, \qquad i < n.$

For $\mathcal{E}_{i,j'}^m$ with $m \in \mathbb{Z}_+$, we have

Lemma 3.10. *For $m \in \mathbb{Z}_+$, the following equalities hold*

(1) $e_{i-1}\mathcal{E}_{i,j'}^m = r^{2m}\mathcal{E}_{i,j'}^m e_{i-1} + [m]\ \mathcal{E}_{i,j'}^{m-1}\mathcal{E}_{i-1,j'}, \qquad i < j \leq n;$

(2) $$e_{j-1}\mathcal{E}_{j-1,j'}^m = (rs)^{2m}\mathcal{E}_{j-1,j'}^m e_{j-1} + s^{2(m-1)}[m]\ \mathcal{E}_{j-1,j'}^{m-1}\times$$
$$\times\Big(s^2\mathcal{E}_{j-1,j+1'}\mathcal{E}_{j-1,j} - s^{-2}\ \mathcal{E}_{j-1,j}\mathcal{E}_{j-1,j+1'}\Big), \qquad j < n;$$

(3) $\mathcal{E}_{i,j'}^m e_{j-1} = s^{-2m}e_{j-1}\mathcal{E}_{i,j'}^m + (rs)^{-2(m-1)}[m]\ \mathcal{E}_{i,j-1'}\mathcal{E}_{i,j'}^{m-1}, \qquad i+1 < j \leq n;$

(4) $\mathcal{E}_{i,j+1'}^m\mathcal{E}_{i,j} \equiv (rs)^{-2m}\Big(\mathcal{E}_{i,j}\mathcal{E}_{i,j+1'}^m + [m]\,\mathcal{E}_{i,j+1}\mathcal{E}_{i,j+2'}\mathcal{E}_{i,j+1'}^{m-1}\Big), \quad i \leq j < n-1;$

(5) $\mathcal{E}_{i,n'}^m\mathcal{E}_{i,n-1} = (rs)^{-2m}\Big(\mathcal{E}_{i,n-1}\mathcal{E}_{i,n'}^m + s^3(r-s)[m]\,\mathcal{E}_{i,n}^2\mathcal{E}_{i,n'}^{m-1}\Big);$

(6) $\mathcal{E}_{i,j+2'}e_{j+1}^m = s^{-2m}e_{j+1}^m\mathcal{E}_{i,j+2'} + (rs)^{2(1-m)}[m]\,e_{j+1}^{m-1}\mathcal{E}_{i,j+1'}, \quad i \leq j < n-1.$

Next, we consider the commutation relations of $\mathcal{E}_{i,j}^m$ with f_i $(1 \leq i \leq n)$.

Lemma 3.11. *For $m \in \mathbb{Z}_+$, the following equalities hold:*

(1) $\mathcal{E}_{i,j}^m f_i = f_i\mathcal{E}_{i,j}^m - [m]\ \mathcal{E}_{i+1,j}\mathcal{E}_{i,j}^{m-1}\omega_i, \qquad i < j < n;$

(2) $\mathcal{E}_{i,j}^m f_j = f_j\mathcal{E}_{i,j}^m + (rs)^{-2(m-1)}[m]\ \mathcal{E}_{i,j-1}\mathcal{E}_{i,j}^{m-1}\omega_j', \qquad i < j < n;$

(3) $e_i^m f_i = f_i e_i^m + (r_i - s_i)^{-1}[m]_i\ e_i^{m-1}(s_i^{-m+1}\omega_i - r_i^{-m+1}\omega_i'), \qquad 1 \leq i \leq n;$

(4) $$\mathcal{E}_{i,n}^m f_i = f_i\mathcal{E}_{i,n}^m - r^{m-1}\beta_m\mathcal{E}_{i+1,n}\mathcal{E}_{i,n}^{m-1}\omega_i - \alpha_m\mathcal{E}_{i+1,n-1}\mathcal{E}_{i,n'}\mathcal{E}_{i,n}^{m-2}\omega_i$$
$$+ r^2\alpha_m\mathcal{E}_{i,n'}\mathcal{E}_{i+1,n-1}\mathcal{E}_{i,n}^{m-2}\omega_i, \qquad i < n-1;$$

(5) $\mathcal{E}_{n-1,n}^m f_{n-1} = f_{n-1}\mathcal{E}_{n-1,n}^m - r^{m-1}\beta_m e_n\mathcal{E}_{n-1,n}^{m-1}\omega_{n-1} - \alpha_m\mathcal{E}_{n-1,n'}\mathcal{E}_{n-1,n}^{m-2}\omega_{n-1};$

(6) $\mathcal{E}_{i,n}^m f_n = f_n\mathcal{E}_{i,n}^m + (r+s)s^{-m-1}\,\beta_m\,\mathcal{E}_{i,n-1}\omega_n'\mathcal{E}_{i,n}^{m-1}, \qquad i < n.$

Lemma 3.12. *For $m \in \mathbb{Z}_+$, the following equalities hold*

(1) $\mathcal{E}_{i,n'}^m f_i = f_i\mathcal{E}_{i,n'}^m - [m]\ \mathcal{E}_{i+1,n'}\mathcal{E}_{i,n'}^{m-1}\omega_i, \qquad i < n-1;$

(2) $\mathcal{E}_{n-1,n'}^m f_{n-1} = f_{n-1}\mathcal{E}_{n-1,n'}^m + (r-s)\,s^{2m-1}[m]\,e_n^2\mathcal{E}_{n-1,n'}^{m-1}\omega_{n-1};$

(3) $\mathcal{E}_{i,n'}^m f_n = f_n\mathcal{E}_{i,n'}^m + (rs)^{-2m+1}(r+s)\,\beta_{2m}\,\mathcal{E}_{i,n}\mathcal{E}_{i,n'}^{m-1}\omega_n', \qquad i \leq n-1;$

(4) $\mathcal{E}_{i,j'}^m f_i = f_i\mathcal{E}_{i,j'}^m - [m]\ \mathcal{E}_{i+1,j'}\mathcal{E}_{i,j'}^{m-1}\omega_i, \qquad i < j-1;$

(5) $$\mathcal{E}_{j-1,j'}^m f_{j-1} = f_{j-1}\mathcal{E}_{j-1,j'}^m + s^{2(m-1)}[m]\times$$
$$\times\big(s^{-2}e_j\mathcal{E}_{j,j+1'} - s^2\mathcal{E}_{j,j+1'}e_j\big)\,\mathcal{E}_{j-1,j'}^{m-1}\omega_{j-1}, \qquad j \leq n-1;$$

(6) $\mathcal{E}_{i,j'}^m f_j = f_j\mathcal{E}_{i,j'}^m + s^{-2}[m]\ \mathcal{E}_{i,j+1'}\mathcal{E}_{i,j'}^{m-1}\omega_j', \qquad i < j < n.$

Proposition 3.13. *$\mathcal{E}_{k,j}^\ell(k \leq j \leq n)$, $\mathcal{E}_{k,m'}^\ell(k < m \leq n)$ commute with f_i for $1 \leq i \leq n$.*

Proof. The proof is carried out in three steps.

Step 1. We want to prove $\mathcal{E}_{k,j}^{\ell}\ \ (k\le j\le n)$ commutes with f_i for $1\le i\le n$. If $i<k$ or $i>j$, this statement is immediate from $(B4)$; while if $i=k$ or $i=j$, it follows directly from Lemma 3.12. We may suppose that $k<i<j$, using $\omega_i'\mathcal{E}_{i+1,j}=r^2\mathcal{E}_{i+1,j}\omega_i'$ and Lemmas 3.1 (3) & 3.12 (2), we obtain

$$\begin{aligned}\mathcal{E}_{k,j}f_i&=(\mathcal{E}_{k,i}\mathcal{E}_{i+1,j}-r^2\mathcal{E}_{i+1,j}\mathcal{E}_{k,i})f_i\\&=(f_i\mathcal{E}_{k,i}+s^{-2}\mathcal{E}_{k,i-1}\omega_i')\mathcal{E}_{i+1,j}-r^2\mathcal{E}_{i+1,j}(f_i\mathcal{E}_{k,i}+s^{-2}\mathcal{E}_{k,i-1}\omega_i')\\&=f_i\mathcal{E}_{k,j}.\end{aligned}$$

Step 2. We want to prove $\mathcal{E}_{k,n'}^{\ell}\ (k<n)$ commutes with f_i for $1\le i\le n$. If $i<k$, this statement is immediate from $(B4)$; while if $i=k$ or $i=n$, it follows directly from Lemma 3.12; if $k<i<n$, it is a similar argument as Step 1.

Step 3. We want to prove $\mathcal{E}_{k,m'}^{\ell}\ (k<m<n)$ commutes with f_i for $1\le i\le n$. If $i<k$, this statement is immediate from $(B4)$; while if $i=k$ or $i=m$, it follows directly from Lemma 3.13. We may suppose that $k\le i\le n$. If $k<i<m$, using $\omega_i'\mathcal{E}_{i+1,m'}=r^2\mathcal{E}_{i+1,m'}\omega_i'$ and Lemma 3.1 (2), we obtain

$$\begin{aligned}\mathcal{E}_{k,m'}f_i&=(\mathcal{E}_{k,i}\mathcal{E}_{i+1,m'}-r^2\mathcal{E}_{i+1,m'}\mathcal{E}_{k,i})f_i\\&=(f_i\mathcal{E}_{k,i}+s^{-2}\mathcal{E}_{k,i-1}\omega_i')\mathcal{E}_{i+1,m'}-r^2\mathcal{E}_{i+1,m'}(f_i\mathcal{E}_{k,i}+s^{-2}\mathcal{E}_{k,i-1}\omega_i')\\&=f_i\mathcal{E}_{k,m'}.\end{aligned}$$

The last case is $m<i\le n$. By Lemma 3.13 (6), we have $[\,\mathcal{E}_{k,i'},f_i]=s^{-2}\mathcal{E}_{k,i+1'}\omega_i'$. Noting that $[\,\mathcal{E}_{k,i+1'}\omega_i',e_{i-1}]_{s^{-2}}=0$, we have

$$\begin{aligned}[\,\mathcal{E}_{k,j'},f_i\,]&=[\,[\cdots[\,\mathcal{E}_{k,i'},e_{i-1}]_{s^{-2}},\cdots,e_j\,]_{s^{-2}},f_i\,]\\&=[\cdots,[\,[\,\mathcal{E}_{k,i'},f_i\,],e_{i-1}\,]_{s^{-2}},\cdots,e_j\,]_{s^{-2}}\\&=0.\end{aligned}$$

This completes the proof. □

Theorem 3.14. *The elements* $\mathcal{E}_{k,j}^{\ell}\ (1\le k\le j\le n)$, $\mathcal{E}_{k,j'}^{\ell}\ (1\le k<j\le n)$, $\mathcal{F}_{k,j}^{\ell}\ (1\le k\le j\le n)$, $\mathcal{F}_{k,j'}^{\ell}\ (1\le k<j\le n)$ *and* $\omega_k^{\ell}-1,\ (\omega_k')^{\ell}-1\ (1\le k\le n)$ *are central in* $U_{r,s}(\mathfrak{so}_{2n+1})$.

Proof. It follows from Propositions 3.11 & 3.14 that $\mathcal{E}_{k,j}^{\ell}\ (1\le k\le j\le n)$, $\mathcal{E}_{k,n'}^{\ell}\ (1\le k<n)$, $\mathcal{E}_{k,j'}^{\ell}\ (1\le k<j\le n-1)$ are central in $U_{r,s}(\mathfrak{so}_{2n+1})$. Applying τ to $\mathcal{E}_{k,j}^{\ell}$ & $\mathcal{E}_{k,j'}^{\ell}$, we see that $\mathcal{F}_{k,j}^{\ell}\ (1\le k\le j\le n)$, $\mathcal{F}_{k,j'}^{\ell}\ (1\le k<j\le n)$ are central. Easy to see that $\omega_k^{\ell}-1,\ (\omega_k')^{\ell}-1\ (1\le k\le n)$ are central, too. □

Remark: If $\ell=2\ell'$, then $\mathcal{E}_{k,j}^{\ell'}\ (1\le k\le j<n)$, $\mathcal{E}_{k,j'}^{\ell'}\ (1\le k<j\le n)$, e_n^{ℓ}, $\mathcal{E}_{k,n}^{\ell}$; $\mathcal{F}_{k,j}^{\ell'}\ (1\le k\le j<n)$, $\mathcal{F}_{k,j'}^{\ell'}\ (1\le k<j\le n)$, f_n^{ℓ}, $\mathcal{F}_{k,n}^{\ell}$; and $\omega_k^{\ell'}-1$, $(\omega_k')^{\ell'}-1\ (1\le k<n)$, $\omega_n^{\ell}-1$, $(\omega_n')^{\ell}-1$ are central in $U_{r,s}(\mathfrak{so}_{2n+1})$.

4. Restricted two-parameter quantum groups

4.1. In what follows, *we assume that ℓ is odd.*

Definition 4.1. The *restricted two-parameter quantum group* is the quotient

$$\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1}):=U_{r,s}(\mathfrak{so}_{2n+1})/I_n,$$

where I_n is the ideal of $U_{r,s}(\mathfrak{so}_{2n+1})$ generated by $\mathcal{E}_{k,j}^{\ell}$ $(1 \leq k \leq j \leq n)$, $\mathcal{E}_{k,j'}^{\ell}$ $(1 \leq k < j \leq n)$, $\mathcal{F}_{k,j}^{\ell}$ $(1 \leq k \leq j \leq n)$, $\mathcal{F}_{k,j'}^{\ell}$ $(1 \leq k < j \leq n)$ and $\omega_k^{\ell} - 1$, $(\omega_k')^{\ell} - 1$ $(1 \leq k \leq n)$.

By Theorem 2.5 and Corollary 2.7, $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ is an algebra of dimension ℓ^{2n^2+2n} with linear basis

$$\mathcal{E}_{1,1}^{a_1}\mathcal{E}_{1,2}^{a_2}\cdots\mathcal{E}_{n-1,n'}^{a_{n^2-1}}\mathcal{E}_{n,n}^{a_{n^2}}\,\omega_1^{b_1}\cdots\omega_n^{b_n}(\omega_1')^{b_1'}\cdots(\omega_n')^{b_n'}\mathcal{F}_{1,1}^{a_1'}\mathcal{F}_{1,2}^{a_2'}\cdots\mathcal{F}_{n-1,n'}^{a'_{n^2-1}}\mathcal{F}_{n,n}^{a'_{n^2}},$$

where all powers range between 0 and $\ell - 1$.

4.2. The main theorem of this section is

THEOREM 4.2. *The ideal I_n is a Hopf ideal, so that $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ is a finite dimensional Hopf algebra.*

4.3. The proof of Theorem 4.2 will be done through a sequence of Lemmas. To determine $\Delta(\mathcal{E}_{i,j}^{\ell})$, $\Delta(\mathcal{E}_{k,j'}^{\ell})$, recall $j' = 2n - j + 1$ given in Section 2. We make conventions:

$$\omega_{i,j} = \prod_{k=i}^{j}\omega_k \ (i \leq j), \quad \omega_{i,j'} = \omega_{i,n}\omega_n\cdots\omega_j \ (i < j), \quad \zeta = s^2 - r^2; \tag{4.1}$$

$$\mathcal{E}_{n+1,n+1} = e_n, \quad \mathcal{E}_{n,n+1} = (s^2 - rs)e_n^2; \tag{4.2}$$

$$\mathcal{E}_{2n-k,2n-k} = r^{-2}s^{-2}e_{k+1}, \quad 1 \leq k < n-1; \tag{4.3}$$

$$\mathcal{E}_{k,2n-k+1} = s^2\mathcal{E}_{k,2n-k}e_k - s^{-2}e_k\mathcal{E}_{k,2n-k}, \ 1 \leq k < n; \tag{4.4}$$

$$\mathcal{E}_{2n-i+1,2n-k+1} = \mathcal{E}_{2n-i+1,2n-k}e_k - s^{-2}e_k\mathcal{E}_{2n-i+1,2n-k}, \ n \geq i > k; \tag{4.5}$$

$$\mathcal{E}_{k+1,2n-k+1} = r^2\mathcal{E}_{k+1,2n-k}e_k - s^{-2}e_k\mathcal{E}_{k+1,2n-k} + \mathcal{E}_{k,2n-k}, \ 1 \leq k < n; \tag{4.6}$$

$$\mathcal{E}_{i,2n-k+1} = \mathcal{E}_{i,2n-k}e_k - s^{-2}e_k\mathcal{E}_{i,2n-k}, \qquad n \geq i > k+1. \tag{4.7}$$

It is easy to verify that

$$\begin{aligned}\mathcal{E}_{k,2n-k+1} = &\sum_{i=0}^{n-k-1}(-1)^i r^{-2(i+1)}s^{-2}\zeta\mathcal{E}_{k,k+i}\mathcal{E}_{k,2n-k-i}\\ &+ (-1)^{n-k}r^{-2(n-k)}(s^2 - rs)\mathcal{E}_{k,n}^2 \in U^+,\end{aligned}$$

$$\mathcal{E}_{2n-i+1,2n-k+1} = \sum_{j=0}^{i-k}(-1)^j r^{-2j}\zeta^{i-k-j}\Big(\sum_b \mathcal{E}_I\Big) \in U^+,$$

where $b = i - k + 1 - j$ and $\mathcal{E}_I = \mathcal{E}_{i_0,i_1}\mathcal{E}_{i_1+1,i_2}\mathcal{E}_{i_2+1,i_3}\cdots\mathcal{E}_{i_{b-1}+1,i_b}$ for $k \leq i_0 \leq i_1 \leq i_2 - 1 \leq \cdots \leq i_b - 1 \leq i - 1$.

A simple calculation shows that $\mathcal{E}_{k+1,2n-k+1} \in U^+$ for $1 \leq k < n$, and $\mathcal{E}_{i,2n-k+1} \in U^+$ for $k + 1 < i \leq n$. So the elements in (4.1)—(4.7) are in U^+.

LEMMA 4.3. (i) $\Delta(\mathcal{E}_{k,j}) = \mathcal{E}_{k,j} \otimes 1 + \omega_{k,j} \otimes \mathcal{E}_{k,j} + \zeta\sum_{i=k}^{j-1}\mathcal{E}_{i+1,j}\omega_{k,i} \otimes \mathcal{E}_{k,i}$ *for* $1 \leq k \leq j \leq n$;

(ii) $\Delta(\mathcal{E}_{k,j'}) = \mathcal{E}_{k,j'} \otimes 1 + \omega_{k,j'} \otimes \mathcal{E}_{k,j'} + \zeta\sum_{i=k}^{2n-j}\mathcal{E}_{i+1,2n-j+1}\omega_{k,i} \otimes \mathcal{E}_{k,i}$ *for* $1 \leq k < j \leq n$.

Proof. (i): The statement is true when $k=j$, owing to the coproduct of e_k. Assume that it is true for $k+1$. Then we obtain

$$\begin{aligned}
\Delta(\mathcal{E}_{k,j}) &= \Delta(e_k)\Delta(\mathcal{E}_{k+1,j}) - r^2\Delta(\mathcal{E}_{k+1,j})\Delta(e_k)\\
&= (e_k\otimes 1+\omega_k\otimes e_k)\Big(\mathcal{E}_{k+1,j}\otimes 1+\omega_{k+1,j}\otimes\mathcal{E}_{k+1,j}+\zeta\sum_{i=k+1}^{j-1}\mathcal{E}_{i+1,j}\omega_{k+1,i}\otimes\mathcal{E}_{k+1,i}\Big)\\
&\quad -r^2\Big(\mathcal{E}_{k+1,j}\otimes 1+\omega_{k+1,j}\otimes\mathcal{E}_{k+1,j}+\zeta\sum_{i=k+1}^{j-1}\mathcal{E}_{i+1,j}\omega_{k+1,i}\otimes\mathcal{E}_{k+1,i}\Big)(e_k\otimes 1+\omega_k\otimes e_k)\\
&= e_k\mathcal{E}_{k+1,j}\otimes 1+\omega_k\mathcal{E}_{k+1,j}\otimes e_k+e_k\omega_{k+1,j}\otimes\mathcal{E}_{k+1,j}+\omega_{k,j}\otimes e_k\mathcal{E}_{k+1,j}\\
&\quad +\zeta\sum_{i=k+1}^{j-1}e_k\mathcal{E}_{i+1,j}\omega_{k+1,i}\otimes\mathcal{E}_{k+1,i}+\zeta\sum_{i=k+1}^{j-1}\omega_k\mathcal{E}_{i+1,j}\omega_{k+1,i}\otimes e_k\mathcal{E}_{k+1,i}\\
&\quad -r^2\mathcal{E}_{k+1,j}e_k\otimes 1-r^2\mathcal{E}_{k+1,j}\omega_k\otimes e_k-r^2\omega_{k+1,j}e_k\otimes\mathcal{E}_{k+1,j}\\
&\quad -r^2\omega_{k+1,j}\omega_k\otimes\mathcal{E}_{k+1,j}e_k-r^2\zeta\sum_{i=k+1}^{j-1}\mathcal{E}_{i+1,j}\omega_{k+1,i}e_k\otimes\mathcal{E}_{k+1,i}\\
&\quad -r^2\zeta\sum_{i=k+1}^{j-1}\mathcal{E}_{i+1,j}\omega_{k+1,i}\omega_k\otimes\mathcal{E}_{k+1,i}e_k\\
&= \mathcal{E}_{k,j}\otimes 1+\omega_{k,j}\otimes\mathcal{E}_{k,j}+\zeta\mathcal{E}_{k+1,j}\omega_k\otimes e_k+\zeta\sum_{i=k+1}^{j-1}\mathcal{E}_{i+1,j}\omega_{k,i}\otimes\mathcal{E}_{k,i}\\
&= \mathcal{E}_{k,j}\otimes 1+\omega_{k,j}\otimes\mathcal{E}_{k,j}+\zeta\sum_{i=k}^{j-1}\mathcal{E}_{i+1,j}\omega_{k,i}\otimes\mathcal{E}_{k,i}
\end{aligned}$$

by using induction hypothesis.

(ii): Use induction on j. For $j=n$, the argument proceeds by induction on k. We have

$$\begin{aligned}
\Delta(\mathcal{E}_{k,n'}) &= \mathcal{E}_{k,n'}\otimes 1+\omega_{k,n'}\otimes\mathcal{E}_{k,n'}+\zeta\sum_{i=k}^{n-2}\mathcal{E}_{i+1,n'}\omega_{k,i}\otimes\mathcal{E}_{k,i}\\
&\quad +(s^2-rs)\zeta e_n^2\omega_{k,n-1}\otimes\mathcal{E}_{k,n-1}+\zeta e_n\omega_{k,n}\otimes\mathcal{E}_{k,n}\\
&= \mathcal{E}_{k,n'}\otimes 1+\omega_{k,n'}\otimes\mathcal{E}_{k,n'}+\zeta\sum_{i=k}^{n}\mathcal{E}_{i+1,n+1}\omega_{k,i}\otimes\mathcal{E}_{k,i}.
\end{aligned}$$

Assume that it is true for $j+1$, we obtain

$$\begin{aligned}
\Delta(\mathcal{E}_{k,j'}) = \Big(&\mathcal{E}_{k,j+1'}\otimes 1+\omega_{k,j+1'}\otimes\mathcal{E}_{k,j+1'}+\zeta\sum_{i=k}^{2n-j-1}\mathcal{E}_{i+1,2n-j}\omega_{k,i}\otimes\mathcal{E}_{k,i}\Big)\\
&\times(e_j\otimes 1+\omega_j\otimes e_j)-s^{-2}(e_j\otimes 1+\omega_j\otimes e_j)\Big(\mathcal{E}_{k,j+1'}\otimes 1\\
&+\omega_{k,j+1'}\otimes\mathcal{E}_{k,j+1'}+\zeta\sum_{i=k}^{2n-j-1}\mathcal{E}_{i+1,2n-j}\omega_{k,i}\otimes\mathcal{E}_{k,i}\Big)
\end{aligned}$$

$$\begin{aligned}
&= \mathcal{E}_{k,j+1'}e_j \otimes 1 + \omega_{k,j+1'}e_j \otimes \mathcal{E}_{k,j+1'} + \zeta \sum_{i=k}^{2n-j-1} \mathcal{E}_{i+1,2n-j}\omega_{k,i}e_j \otimes \mathcal{E}_{k,i}\\
&\quad + \mathcal{E}_{k,j+1'}\omega_j \otimes e_j + \omega_{k,j+1'}\omega_j \otimes \mathcal{E}_{k,j+1'}e_j\\
&\quad + \zeta \sum_{i=k}^{2n-j-1} \mathcal{E}_{i+1,2n-j}\omega_{k,i}\omega_j \otimes \mathcal{E}_{k,i}e_j\\
&\quad - s^{-2}e_j\mathcal{E}_{k,j+1'} \otimes 1 - s^{-2}e_j\omega_{k,j+1'} \otimes \mathcal{E}_{k,j+1'}\\
&\quad - s^{-2}\zeta \sum_{i=k}^{2n-j-1} e_j\mathcal{E}_{i+1,2n-j}\omega_{k,i} \otimes \mathcal{E}_{k,i}\\
&\quad - s^{-2}\omega_j\mathcal{E}_{k,j+1'} \otimes e_j - s^{-2}\omega_j\omega_{k,j+1'} \otimes e_j\mathcal{E}_{k,j+1'}\\
&\quad - s^{-2}\zeta \sum_{i=k}^{2n-j-1} \omega_j\mathcal{E}_{i+1,2n-j}\omega_{k,i} \otimes e_j\mathcal{E}_{k,i}\\
&= \mathcal{E}_{k,j'} \otimes 1 + \omega_{k,j'} \otimes \mathcal{E}_{k,j'} + \zeta\mathcal{E}_{2n-j+1,2n-j+1}\omega_{k,2n-j} \otimes \mathcal{E}_{k,2n-j}\\
&\quad + \zeta \sum_{i=k}^{j-2} \mathcal{E}_{i+1,2n-j+1}\omega_{k,i} \otimes \mathcal{E}_{k,i} + \zeta\mathcal{E}_{j,2n-j+1}\omega_{k,j-1} \otimes \mathcal{E}_{k,j-1}\\
&\quad + \zeta\mathcal{E}_{j+1,2n-j+1}\omega_{k,j} \otimes \mathcal{E}_{k,j} + \zeta \sum_{i=j+1}^{n} \mathcal{E}_{i+1,2n-j+1}\omega_{k,i} \otimes \mathcal{E}_{k,i}\\
&\quad + \zeta \sum_{i=n+1}^{2n-j-1} \mathcal{E}_{i+1,2n-j+1}\omega_{k,i} \otimes \mathcal{E}_{k,i}\\
&= \mathcal{E}_{k,j'} \otimes 1 + \omega_{k,j'} \otimes \mathcal{E}_{k,j'} + \zeta \sum_{i=k}^{2n-j} \mathcal{E}_{i+1,2n-j+1}\omega_{k,i} \otimes \mathcal{E}_{k,i}
\end{aligned}$$

by induction hypothesis and notations in (4.1)—(4.7) and Lemma 3.2 (3). □

4.4. We generalize Lemma 4.3 (i) to an expression for $\Delta(\mathcal{E}_{k,j}^a)$ in Lemma 4.5 below. The formula involves certain exponents p_m of s that are defined recursively on ordered m-tuples of nonnegative integers, as follows (see [**BW3**]):

$$\begin{aligned}
p_m(0, \cdots, 0) &:= 0,\\
p_m(c_1+1, c_2, \cdots, c_m) &:= p_m(c_1, c_2, \cdots, c_m) - c_2 - c_3 - \cdots - c_m,\\
p_m(c_1, c_2, \cdots, c_{m-1}, c_m+1) &:= p_m(c_1, c_2, \cdots, c_m) - c_1 - c_2 - \cdots - c_{m-1},\\
p_m(c_1, c_2, \cdots, c_j+1, \cdots, c_m) &:= p_m(c_1, c_2, \cdots, c_m) - c_1 - c_2 - \cdots - c_{j-1}\\
&\qquad + c_j - c_{j+1} - \cdots - c_m, \quad (1 < j < m).
\end{aligned}$$

An inductive argument on $c_1 + \cdots + c_m$ shows that p_m is well-defined.

Set $C_m^a(r,s) = \frac{[c_1+\cdots+c_m]!}{[c_1]!\cdots[c_m]!}$, for $a = c_1 + \cdots + c_m$. The lemma below is true, which is equivalent to checking inductively: $[\,c_1 + \cdots + c_m\,] = \sum_{j=1}^{m} r^{2C_j}s^{2D_j}[\,c_j\,]$.

Lemma 4.4. *Let $c_1, \cdots, c_m$ be positive integers with $a = c_1 + \cdots + c_m$. Then*

$$C_m^a(r,s) = \sum_{j=1}^{m} r^{2C_j}s^{2D_j} \begin{bmatrix} c_1 + \cdots + c_m - 1 \\ c_m \end{bmatrix} \cdots \begin{bmatrix} c_1 + \cdots + c_j - 1 \\ c_j - 1 \end{bmatrix} \cdots \begin{bmatrix} c_1 + c_2 \\ c_2 \end{bmatrix},$$

where $C_j = c_{j+1} + c_{j+2} + \cdots + c_m, D_j = c_1 + \cdots + c_{j-1}$, and $C_m = 0 = D_1$.

In the above sum, the binomial coefficients have 1 subtracted from their top number to the one with $c_1 + \cdots + c_j$ on top and not thereafter, as in the jth summand we have replaced c_j by $c_j - 1$.

LEMMA 4.5. *For* $1 \le k \le j \le n$ *and* $m = j - k + 2$. *Then*

$$\begin{aligned}\Delta(\mathcal{E}_{k,j}^{a}) = \sum s^{2p_m(c_1,\cdots,c_m)} \zeta^{a-c_1-c_m} C_m^a(r,s) \mathcal{E}_{k,j}^{c_1} \mathcal{E}_{k+1,j}^{c_2} \cdots \mathcal{E}_{j,j}^{c_{m-1}} \omega_{k,k}^{c_2} \omega_{k,k+1}^{c_3} \cdots \omega_{k,j}^{c_m} \\ \otimes \mathcal{E}_{k,k}^{c_2} \mathcal{E}_{k,k+1}^{c_3} \cdots \mathcal{E}_{k,j}^{c_m},\end{aligned}$$

the sum taken over all m*-tuples* $(c_1, \cdots, c_m)$ *with* $c_1 + \cdots + c_m = a$.

PROOF. Observe first that if $a = 1$, the above expression coincides with that in Lemma 4.3 (i). To simplify notation, we will assume without loss of generality that $k = 1$ (and hence $m = j + 1$). Now suppose that the above formula holds for $a - 1$, that is $\Delta(\mathcal{E}_{1,j}^{a-1}) = \sum s^{2p_{j+1}(c_1,\cdots,c_{j+1})} \zeta^{a-1-c_1-c_{j+1}} C_{j+1}^{a-1}(r,s) \mathcal{E}_{1,j}^{c_1} \cdots \mathcal{E}_{j,j}^{c_j} \omega_{1,1}^{c_2} \cdots \omega_{1,j}^{c_{j+1}}$ $\otimes \mathcal{E}_{1,1}^{c_2} \cdots \mathcal{E}_{1,j}^{c_{j+1}}$, where the sum is over all $(j{+}1)$-tuples $(c_1, \cdots, c_{j+1})$ with $c_1 + \cdots + c_{j+1} = a - 1$. Then $\Delta(\mathcal{E}_{1,j}^{a}) = \Delta(\mathcal{E}_{1,j}^{a-1})\Delta(\mathcal{E}_{1,j})$, which by Lemma 4.3 (i) is equal to the above sum times

$$\mathcal{E}_{1,j} \otimes 1 + \omega_{1,j} \otimes \mathcal{E}_{1,j} + \zeta \sum_{i=1}^{j-1} \mathcal{E}_{i+1,j} \omega_{1,i} \otimes \mathcal{E}_{1,i}.$$

Expanding and re-summing over all $(d_1, \cdots, d_{j+1})$ with $d_1 + \cdots + d_{j+1} = a$, by Lemma 3.3 (4) & (5), we obtain

$$\begin{aligned}&\textit{the coefficient of } \mathcal{E}_{1,j}^{d_1} \mathcal{E}_{2,j}^{d_2} \cdots \mathcal{E}_{j,j}^{d_j} \omega_{1,1}^{d_2} \omega_{1,2}^{d_3} \cdots \omega_{1,j}^{d_{j+1}} \otimes \mathcal{E}_{1,1}^{d_2} \mathcal{E}_{1,2}^{d_3} \cdots \mathcal{E}_{1,j}^{d_{j+1}} \\ &\quad = s^{2p_{j+1}(d_1,\cdots,d_{j+1})} \zeta^{a-d_1-d_{j+1}} \times \\ &\quad\quad \times \Big(\sum_{i=1}^{j+1} r^{2C_i'} s^{2D_i'} \begin{bmatrix} d_1 + \cdots + d_{j+1} - 1 \\ d_{j+1} \end{bmatrix} \cdots \begin{bmatrix} d_1 + \cdots + d_i - 1 \\ d_i - 1 \end{bmatrix} \cdots \begin{bmatrix} d_1 + d_2 \\ d_2 \end{bmatrix} \Big),\end{aligned}$$

where $C_i' = d_{i+1} + d_{i+2} + \cdots + d_{j+1}$, $D_i' = d_1 + \cdots + d_{i-1}$ and $C_{j+1}' = 0 = D_1'$. By Lemma 4.4, we get the desired result. □

As a result, we have the following formula for $e_j = \mathcal{E}_{j,j}$ for $1 \le j \le n$:

$$\Delta(e_j^a) = \sum_{i=0}^{a} s^{2i(i-a)} \begin{bmatrix} a \\ i \end{bmatrix} e_j^i \omega_j^{a-i} \otimes e_j^{a-i}. \tag{4.8}$$

4.5. We generalize Lemma 4.5 to a formula for $\Delta(\mathcal{E}_{k,n'}^{\ell})$ in Lemma 4.6 below. The formula involves certain exponents $p_m (m = n - k + 3)$ of s that are defined recursively on ordered m-tuples of nonnegative integers, as follows (which are different from those in Lemma 4.5):

$$\begin{aligned} p_m(0, \cdots, 0) &:= 0, \\ p_m(c_1 + 1, c_2, \cdots, c_m) &:= p_m(c_1, c_2, \cdots, c_m) - c_2 - c_3 - \cdots - c_m, \\ p_m(c_1, c_2, \cdots, c_{m-1}, c_m + 1) &:= p_m(c_1, c_2, \cdots, c_m) - c_1 - c_2 - \cdots - c_{m-1},\end{aligned}$$

$$\begin{aligned} p_m(c_1,c_2,\cdots,c_j+1,\cdots,c_m) &:= p_m(c_1,c_2,\cdots,c_m)-c_1-c_2-\cdots-c_{j-1}\\ &\quad +c_j-c_{j+1}-\cdots-c_m, \quad (1<j<m-2),\\ p_m(c_1,c_2,\cdots,c_{m-2}+1,c_{m-1},c_m) &:= p_m(c_1,c_2,\cdots,c_m)-c_1-\cdots-c_{m-3}\\ &\quad +2c_{m-2}-c_m,\\ p_m(c_1,c_2,\cdots,c_{m-2},c_{m-1}+1,c_m) &:= p_m(c_1,c_2,\cdots,c_m)-c_1-\cdots-c_{m-3}-c_m. \end{aligned}$$

An inductive argument on $c_1+\cdots+c_m$ shows that p_m is well-defined.

Lemma 4.6. *For $1\le k<n$ and $m=n-k+3$, we have*

$$\begin{aligned} \Delta(\mathcal{E}^a_{k,n'}) = &\sum_{c_{m-1}=0,1} s^{2p_m(c_1,\cdots,c_m)}\zeta^{a-c_1-c_m}C^a_m(r,s)\mathbf{E}^{(a)}_k\\ &+\zeta(rs-r^2)[a]!\sum_{2\le c_{m-1}<a} s^{2p_m(c_1,\cdots,c_m)}\mathbf{E}^{(a)}_k\\ &+r^{\frac{a(a-1)}{2}}(s-r)^{a-1}\zeta\Big(\prod_{i=2}^{a}(r^i+s^i)\Big)e^a_n\omega^a_{k,n}\otimes\mathcal{E}^a_{k,n}, \end{aligned}$$

where $\mathbf{E}^{(a)}_k=\mathcal{E}^{c_1}_{k,n+1}\cdots\mathcal{E}^{c_{n-k+2}}_{n+1,n+1}\omega^{c_2}_{k,k}\cdots\omega^{c_j}_{k,k+j-2}\cdots\omega^{c_{n-k+3}}_{k,n+1}\otimes\mathcal{E}^{c_2}_{k,k}\cdots\mathcal{E}^{c_j}_{k,k+j-2}\cdots\times$ $\times\,\mathcal{E}^{c_{n-k+3}}_{k,n+1}$ *with* $c_1+\cdots+c_m=a$.

Proof. Observe first that if $a=1$, the above expression only has the first term which coincides with that in Lemma 4.3 (ii). To simplify notation, we will assume without loss of generality that $k=1$ (and hence $m=n+2$). Now suppose that the above formula holds for $a-1$, that is,

$$\begin{aligned} \Delta(\mathcal{E}^{a-1}_{1,n'}) = &\sum_{\substack{c_{n+1}=0,1\\ c_1+\cdots+c_{n+2}=a-1}} s^{2p_{n+2}(c_1,\cdots,c_{n+2})}\zeta^{a-1-c_1-c_{n+2}}C^{a-1}_{n+2}(r,s)\mathbf{E}^{(a-1)}_1\\ &+\zeta(rs-r^2)[a-1]!\sum_{\substack{2\le c_{n+1}<a-1,\\ c_1+\cdots+c_{n+2}=a-1}} s^{2p_{n+2}(c_1,\cdots,c_{n+2})}\mathbf{E}^{(a-1)}_1\\ &+r^{\frac{(a-2)(a-1)}{2}}(s-r)^{a-2}\zeta\Big(\prod_{i=2}^{a-1}(r^i+s^i)\Big)e^{a-1}_n\omega^{a-1}_{k,n}\otimes\mathcal{E}^{a-1}_{k,n}, \end{aligned}$$

where the sum is over all $(n+2)$-tuples $(c_1,\cdots,c_{n+2})$ with $c_1+\cdots+c_{n+2}=a-1$. Then $\Delta(\mathcal{E}^a_{1,n'})=\Delta(\mathcal{E}^{a-1}_{1,n'})\Delta(\mathcal{E}_{1,n'})$, which by Lemma 4.3 (ii) is equal to the above sum times

$$\mathcal{E}_{1,n'}\otimes 1+\omega_{1,n'}\otimes\mathcal{E}_{1,n'}+\zeta\sum_{i=1}^{n}\mathcal{E}_{i+1,n'}\omega_{1,i}\otimes\mathcal{E}_{1,i}.$$

Expanding and re-summing over all $(d_1,\cdots,d_{n+2})$ with $d_1+\cdots+d_{n+2}=a$, we get the desired result. □

4.6. Proof of Theorem 4.2. We are now in the position to prove that I_n is a Hopf ideal. As $[\ell]=0$, it follows from Lemma 4.5 that the only nonzero terms of $\Delta(\mathcal{E}^\ell_{k,j})$ are those having $c_i=\ell$ for some i. As a result,

$$\Delta(\mathcal{E}^\ell_{k,j})=\mathcal{E}^\ell_{k,j}\otimes 1+\omega^\ell_{k,j}\otimes\mathcal{E}^\ell_{k,j}+s^{\ell(\ell-1)}\zeta^\ell\sum_{i=k}^{j-1}\mathcal{E}^\ell_{i+1,j}\omega^\ell_{k,i}\otimes\mathcal{E}^\ell_{k,i}, \tag{4.9}$$

which is clearly in $I_n \otimes U + U \otimes I_n$. Applying the antipode property to (4.9), we obtain

$$0 = \varepsilon(\mathcal{E}^{\ell}_{k,j}) = \mathcal{E}^{\ell}_{k,j} + \omega^{\ell}_{k,j} S(\mathcal{E}^{\ell}_{k,j}) + s^{\ell(\ell-1)}\zeta^{\ell} \sum_{i=k}^{j-1} \mathcal{E}^{\ell}_{i+1,j}\omega^{\ell}_{k,i} S(\mathcal{E}^{\ell}_{k,i}).$$

So we have

$$S(\mathcal{E}^{\ell}_{k,j}) = -\omega^{-\ell}_{k,j}\big(\mathcal{E}^{\ell}_{k,j} + s^{\ell(\ell-1)}\zeta^{\ell} \sum_{i=k}^{j-1} \mathcal{E}^{\ell}_{i+1,j}\omega^{\ell}_{k,i} S(\mathcal{E}^{\ell}_{k,i})\big). \tag{4.10}$$

If $j = k$, we have $S(\mathcal{E}^{\ell}_{k,j}) = S(e^{\ell}_k) = (-\omega_k^{-1} e_k)^{\ell}$, which is a scalar multiple of $\omega_k^{-\ell} e^{\ell}_k \in I_n$. Using (4.10) and induction on $j - k$, we see that $S(\mathcal{E}^{\ell}_{k,j}) \in I_n$ for all $k < j$ as well. From Lemma 4.6, we have

$$\begin{aligned}
\Delta(\mathcal{E}^{\ell}_{k,n'}) = {} & \mathcal{E}^{\ell}_{k,n'} \otimes 1 + \omega^{\ell}_{k,n'} \otimes \mathcal{E}^{\ell}_{k,n'} + s^{2\ell(\ell-1)}\zeta^{\ell} \mathcal{E}^{\ell}_{n,n'}\omega^{\ell}_{k,n-1} \otimes \mathcal{E}^{\ell}_{k,n-1} \\
& + r^{\frac{\ell(\ell-1)}{2}}(s-r)^{\ell-1}\zeta\Big(\prod_{i=2}^{\ell}(r^i + s^i)\Big) e^{\ell}_n \omega^{\ell}_{k,n} \otimes \mathcal{E}^{\ell}_{k,n} \\
& + s^{\ell(\ell-1)}\zeta^{\ell} \sum_{i=k}^{n-2} \mathcal{E}^{\ell}_{i+1,n'}\omega^{\ell}_{k,i} \otimes \mathcal{E}^{\ell}_{k,i},
\end{aligned} \tag{4.11}$$

which is clearly in $I_n \otimes U + U \otimes I_n$. Applying the antipode property to (4.11), we have $S(\mathcal{E}^{\ell}_{k,n'}) \in I_n$. For $j < n$, using a similar method of Lemma 4.6, we have

$$\begin{aligned}
\Delta(\mathcal{E}^{\ell}_{k,j'}) = {} & \mathcal{E}^{\ell}_{k,j'} \otimes 1 + \omega^{\ell}_{k,j'} \otimes \mathcal{E}^{\ell}_{k,j'} + s^{2\ell(\ell-1)}\zeta^{\ell} \mathcal{E}^{\ell}_{j,j'}\omega^{\ell}_{k,j-1} \otimes \mathcal{E}^{\ell}_{k,j-1} \\
& + r^{\frac{\ell(\ell-1)}{2}}(s-r)^{\ell-1}\zeta\Big(\prod_{i=2}^{\ell}(r^i + s^i)\Big) \mathcal{E}^{\ell}_{n+1,j'} \omega^{\ell}_{k,n} \otimes \mathcal{E}^{\ell}_{k,n} \\
& + s^{\ell(\ell-1)}\zeta^{\ell} \sum_{i=k}^{j-2} \mathcal{E}^{\ell}_{i+1,j'}\omega^{\ell}_{k,i} \otimes \mathcal{E}^{\ell}_{k,i} \\
& + r^{\ell(\ell-1)} s^{2\ell(\ell-1)}\zeta^{\ell} \sum_{i=j}^{n-1} \mathcal{E}^{\ell}_{i+1,j'}\omega^{\ell}_{k,i} \otimes \mathcal{E}^{\ell}_{k,i} \\
& + r^{-\ell(\ell-1)}\zeta^{\ell} \sum_{i=n+1}^{2n-j} \mathcal{E}^{\ell}_{i+1,j'}\omega^{\ell}_{k,i} \otimes \mathcal{E}^{\ell}_{k,i},
\end{aligned} \tag{4.12}$$

which is in $I_n \otimes U + U \otimes I_n$. Applying the antipode property to this, we have $S(\mathcal{E}^{\ell}_{k,j'}) \in I_n$.

Using τ, we find that $\Delta(\mathcal{F}_{k,j}),\ \Delta(\mathcal{F}_{k,j'}) \in I_n \otimes U + U \otimes I_n$. So I_n is a Hopf ideal, and $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ is a finite-dimensional Hopf algebra. $\square$

5. Isomorphisms of $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$, as well as $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$

5.1. Write $\mathfrak{u}_{r,s}=\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$. Let G denote the group generated by ω_i, ω_i' $(1\le i\le n)$ in $\mathfrak{u}=\mathfrak{u}_{r,s}$. Define linear subspace $\mathfrak{a}_k$ of $\mathfrak{u}$ by

$$\begin{aligned}\mathfrak{a}_0=\mathbb{K}G,\qquad \mathfrak{a}_1&=\mathbb{K}G+\sum_{i=1}^n(\mathbb{K}e_iG+\mathbb{K}f_iG),\\ \mathfrak{a}_k&=(\mathfrak{a}_1)^k\quad\text{for}\quad k\ge 1.\end{aligned}\tag{5.1}$$

Note that $1\in\mathfrak{a}_0$, $\Delta(\mathfrak{a}_0)\subseteq\mathfrak{a}_0\otimes\mathfrak{a}_0$, $\mathfrak{a}_1$ generates $\mathfrak{u}$ as an algebra, and $\Delta(\mathfrak{a}_1)\subseteq\mathfrak{a}_1\otimes\mathfrak{a}_0+\mathfrak{a}_0\otimes\mathfrak{a}_1$. By [**M**], $\{\mathfrak{a}_k\}$ is a coalgebra filtration of $\mathfrak{u}$ and $\mathfrak{u}_0\subseteq\mathfrak{a}_0$, where the coradical $\mathfrak{u}_0$ of $\mathfrak{u}$ is the sum of all the simple subcoalgebras of $\mathfrak{u}$. Clearly, $\mathfrak{a}_0\subseteq\mathfrak{u}_0$ as well, and so $\mathfrak{u}_0=\mathbb{K}G$. This implies that $\mathfrak{u}$ is *pointed*, that is, every simple subcoalgebra of $\mathfrak{u}$ is one-dimensional.

Let $\mathfrak{b}$ (resp., $\mathfrak{b}'$) be the Hopf subalgebra of $\mathfrak{u}=\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ generated by $e_i,\omega_i^{\pm1}$ (resp., $f_i,(\omega_i')^{\pm1}$), $1\le i\le n$. The same argument shows that $\mathfrak{b}$ and $\mathfrak{b}'$ are pointed as well. Then we have

Proposition 5.1. *$\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ is a pointed Hopf algebra, as are $\mathfrak{b}$ and $\mathfrak{b}'$.* □

5.2. It follows from [**M**, Lemma 5.5.1] that $\mathfrak{a}_k\subseteq\mathfrak{u}_k$ for all k, where $\{\mathfrak{u}_k\}$ is the *coradical filtration* of $\mathfrak{u}$ defined inductively by $\mathfrak{u}_k=\Delta^{-1}(\mathfrak{u}\otimes\mathfrak{u}_{k-1}+\mathfrak{u}_0\otimes\mathfrak{u})$. In particular, $\mathfrak{a}_1\subseteq\mathfrak{u}_1$. By [**M**, Theorem 5.4.1], as $\mathfrak{u}$ is pointed, $\mathfrak{u}_1$ is spanned by the set of group-like elements G, together with all the skew-primitive elements of $\mathfrak{u}$.

Let m be the smallest positive integer such that $r^m=s^m$. Note that $[m]=0$ while $[1],[2],\cdots,[m-1]$ are all nonzero. It follows from this observation and (4.8) that e_i^m,f_i^m $(1\le k\le n)$ are skew-primitive if $m<\ell$. From now on, we assume that $m=\ell$. Based on Lemmas 4.3, 4.5, 4.6 & formula (4.12), together with with the same argument of [BW3, Lemma 3.3], we have

Lemma 5.2. *Assume that rs^{-1} is a primitive ℓth root of unity. Then*

$$\mathfrak{u}_1=\mathbb{K}G+\sum_{i=1}^n(\mathbb{K}e_iG+\mathbb{K}f_iG).$$

5.3. Given two group-like elements g and h in a Hopf algebra H, let $P_{g,h}(H)$ denote the set of skew-primitive elements of H given by

$$P_{g,h}(H)=\{x\in H\mid\Delta(x)=x\otimes g+h\otimes x\}.$$

Lemma 5.3. *Assume that rs^{-1} is a primitive ℓth root of unity. Then*

(i) $P_{1,\omega_i}(\mathfrak{u}_{r,s})=\mathbb{K}(1-\omega_i)+\mathbb{K}e_i;\quad P_{1,\omega_i'^{-1}}(\mathfrak{u}_{r,s})=\mathbb{K}(1-\omega_i'^{-1})+\mathbb{K}f_i\omega_i'^{-1};$
$P_{1,\sigma}(\mathfrak{u}_{r,s})=\mathbb{K}(1-\sigma),$ *for* $\sigma\notin\{\omega_i,\omega_i'^{-1}\mid1\le i\le n\}$.

(ii) $P_{\omega_i',1}(\mathfrak{u}_{r,s})=\mathbb{K}(1-\omega_i')+\mathbb{K}f_i;\quad P_{\omega_i^{-1},1}(\mathfrak{u}_{r,s})=\mathbb{K}(1-\omega_i^{-1})+\mathbb{K}e_i\omega_i^{-1};$
$P_{\sigma,1}(\mathfrak{u}_{r,s})=\mathbb{K}(1-\sigma),$ *for* $\sigma\notin\{\omega_i^{-1},\omega_i'\mid1\le i\le n\}$.

Proof. Similar to that of Lemma 4.3 in [**HW**]. □

5.4. Lemma 5.3 is crucial for determining the Hopf isomorphisms of $\mathfrak{u}_{r,s}$.

Theorem 5.4. *Assume that rs^{-1} and $r's'^{-1}$ are primitive ℓth roots of unity with $\ell\ne3,4$, and ζ is a 2nd root of unity. Then $\varphi:\mathfrak{u}_{r,s}\cong\mathfrak{u}_{r',s'}$ as Hopf algebras if and only if either*

(1) $(r', s') = \zeta(r, s)$, φ *is a diagonal isomorphism*: $\varphi(\omega_i) = \tilde{\omega}_i$, $\varphi(\omega'_i) = \tilde{\omega}'_i$, $\varphi(e_i) = a_i\tilde{e}_i$, $\varphi(f_i) = \zeta^{\delta_{i,n}} a_i^{-1}\tilde{f}_i$*; or*

(2) $(r', s') = \zeta(s, r)$, $\varphi(\omega_i) = \tilde{\omega}'^{-1}_i$, $\varphi(\omega'_i) = \tilde{\omega}_i^{-1}$, $\varphi(e_i) = a_i\tilde{f}_i\tilde{\omega}'^{-1}_i$, $\varphi(f_i) = \zeta^{\delta_{i,n}} a_i^{-1}\tilde{\omega}_i^{-1}\tilde{e}_i$, *where* $a_i \in \mathbb{K}^*$.

Proof. Suppose $\varphi : \mathfrak{u}_{r,s} \longrightarrow \mathfrak{u}_{r',s'}$ is a Hopf algebra isomorphism. Write $\tilde{e}_i$, $\tilde{f}_i$, $\tilde{\omega}_i$, $\tilde{\omega}'_i$ to distinguish the generators of $\mathfrak{u}_{r',s'}$. Because

$$\Delta(\varphi(e_i)) = (\varphi \otimes \varphi)(\Delta(e_i)) = \varphi(e_i) \otimes 1 + \varphi(\omega_i) \otimes \varphi(e_i),$$

$\varphi(e_i) \in P_{1,\varphi(\omega_i)}(\mathfrak{u}_{r',s'})$. As φ is an isomorphism, $\varphi(\omega_i) \in \mathbb{K}\tilde{G}$. Lemma 5.3 (i) implies $\varphi(\omega_i) \in \{\, \tilde{\omega}_j, \tilde{\omega}'^{-1}_j \mid 1 \leq j \leq n \,\}$. More precisely, we have either $\varphi(\omega_i) = \tilde{\omega}_j$, $\varphi(e_i) = a(1 - \tilde{\omega}_j) + b\tilde{e}_j$, or $\varphi(\omega_i) = \tilde{\omega}'^{-1}_j$, $\varphi(e_j) = a(1 - \tilde{\omega}'^{-1}_j) + b\tilde{f}_j\tilde{\omega}'^{-1}_j$ for some j with $1 \leq j \leq n$ and $a, b \in \mathbb{K}$. But in both cases, we have $a = 0$. The reasoning is the following: Applying φ to relation $\omega_i e_i = r_i s_i^{-1} e_i \omega_i$, we get

$$\tilde{\omega}_j \big[\, a(1 - \tilde{\omega}_j) + b\tilde{e}_j \,\big] = r_i s_i^{-1} \big[\, a(1 - \tilde{\omega}_j) + b\tilde{e}_j \,\big]\tilde{\omega}_j, \quad \textit{or}$$
$$\tilde{\omega}'^{-1}_j \big[\, a(1 - \tilde{\omega}'^{-1}_j) + b\tilde{f}_j\tilde{\omega}'^{-1}_j \,\big] = r_i s_i^{-1} \big[\, a(1 - \tilde{\omega}'^{-1}_j) + b\tilde{f}_j\tilde{\omega}'^{-1}_j \,\big]\tilde{\omega}'^{-1}_j.$$

Since $r_i \neq s_i$, $a = 0$. Thus, $b \in \mathbb{K}^*$.

(I) We claim $\varphi(\omega_n) \in \{\tilde{\omega}_n, \tilde{\omega}'^{-1}_n\}$. Indeed, let us consider the restriction $\varphi|_\bullet$ of φ to the Hopf subalgebra generated by $e_i, f_i, \omega_i^{\pm 1}, \omega_i'^{\pm 1}$ for $i = n-1, n$, which is isomorphic to $\mathfrak{u}_{r,s}(\mathfrak{so}_5)$. According to the above discussion, we have 4 possibilities to occur, namely, case (i): For $i = n{-}1, n$, we have $\varphi(\omega_i) = \tilde{\omega}_{j_i}$ and $\varphi(e_i) = a_i\tilde{e}_{j_i}$ ($j_i \in \{n{-}1, n\}$, $a_i \in \mathbb{K}^*$); case (ii): For $i = n{-}1, n$, we have $\varphi(\omega_i) = \tilde{\omega}'^{-1}_{j_i}$ and $\varphi(e_i) = a_i\tilde{f}_{j_i}\tilde{\omega}'^{-1}_{j_i}$ ($j_i \in \{n{-}1, n\}$, $a_i \in \mathbb{K}^*$); case (iii): $\varphi(\omega_{n-1}) = \tilde{\omega}_n$, but $\varphi(\omega_n) = \tilde{\omega}'^{-1}_{n-1}$; case (iv): $\varphi(\omega_{n-1}) = \tilde{\omega}'^{-1}_n$, but $\varphi(\omega_n) = \tilde{\omega}_{n-1}$.

Case (i): In this case, we claim $\varphi(\omega_i) = \tilde{\omega}_i$ for $i = n{-}1, n$. Otherwise, if $\varphi(\omega_i) = \tilde{\omega}_j$ for $i \neq j \in \{n{-}1, n\}$, then applying φ to relation $\omega_i e_i = r_i s_i^{-1} e_i \omega_i$ yields $\tilde{\omega}_j\tilde{e}_j = r_i s_i^{-1}\tilde{e}_j\tilde{\omega}_j$, it follows that $r'_j s'^{-1}_j = r_i s_i^{-1}$ for $j \neq i \in \{n{-}1, n\}$. This means $(rs^{-1})^3 = 1$, which contradicts the assumption. So we get $\varphi(\omega_i) = \tilde{\omega}_i$ for $i = n{-}1, n$. Similarly, we have $\varphi(\omega'_i) = \tilde{\omega}'_i$ for $i = n{-}1, n$, as φ preserves $(B4)$. Thereby, we get $\varphi(e_i) = a_i\tilde{e}_i$, $\varphi(f_i) = \zeta^{\delta_{i,n}} a_i^{-1}\tilde{f}_i$ for $i = n{-}1, n$, where $a_i \in \mathbb{K}^*$.

Furthermore, applying φ to relations $\omega_i e_j = \langle \omega'_j, \omega_i \rangle e_j \omega_i$ for $i, j \in \{n{-}1, n\}$, we get $\langle \omega'_j, \omega_i \rangle = \langle \tilde{\omega}'_j, \tilde{\omega}_i \rangle$ for $i, j \in \{n{-}1, n\}$, i.e., $r'^2 s'^{-2} = r^2 s^{-2}$, $r' s'^{-1} = rs^{-1}$, $r'^{-2} = r^{-2}$ and $s'^2 = s^2$. Hence, $(r', s') = \zeta(r, s)$, $\varphi|_\bullet$ is a diagonal isomorphism.

Case (ii): In this case, we claim $\varphi(\omega_i) = \tilde{\omega}'^{-1}_i$ for $i = n{-}1, n$. Otherwise, if $\varphi(\omega_i) = \tilde{\omega}'^{-1}_j$ for $i \neq j \in \{n{-}1, n\}$, then $\varphi(e_i) = a_i\tilde{f}_j\tilde{\omega}'^{-1}_j$ for $i \neq j \in \{n{-}1, n\}$. Using φ to relations $\omega_i e_i = \langle \omega'_i, \omega_i \rangle e_i \omega_i$ for $i \in \{n{-}1, n\}$, we get $\tilde{\omega}'^{-1}_j\tilde{f}_j\tilde{\omega}'^{-1}_j = \langle \omega'_i, \omega_i \rangle \tilde{f}_j\tilde{\omega}'^{-1}_j\tilde{\omega}'^{-1}_j$ so that $\langle \tilde{\omega}'_j, \tilde{\omega}_j \rangle^{-1} = \langle \omega'_i, \omega_i \rangle$ for $i \neq j \in \{n{-}1, n\}$. This means $(rs^{-1})^3 = 1$. This contradicts the assumption. So we have $\varphi(\omega_i) = \tilde{\omega}'^{-1}_i$ for $i = n{-}1, n$. Similarly, $\varphi(\omega'_i) = \tilde{\omega}_i^{-1}$ for $i = n{-}1, n$, as φ preserves $(B4)$. Hence, $\varphi(\omega_i) = \tilde{\omega}'^{-1}_i$, $\varphi(\omega'_i) = \tilde{\omega}_i^{-1}$, $\varphi(e_i) = a_i\tilde{f}_i\tilde{\omega}'^{-1}_i$ for $i = n{-}1, n$, where $a_i \in \mathbb{K}^*$.

Using φ to relations $\omega_i e_j = \langle \omega'_j, \omega_i \rangle e_j \omega_i$ for $i, j \in \{n{-}1, n\}$, we get $\langle \tilde{\omega}'_i, \tilde{\omega}_j \rangle^{-1} = \langle \omega'_j, \omega_i \rangle$ for $i, j \in \{n{-}1, n\}$, i.e., $r'^{-2}s'^2 = r^2 s^{-2}$, $(r's'^{-1})^{-1} = rs^{-1}$, $r'^2 = s^2$ and $s'^{-2} = r^{-2}$. Hence, we have $(r', s') = \zeta(s, r)$, $\varphi(f_i) = \zeta^{\delta_{i,n}} a_i^{-1}\tilde{\omega}_i^{-1}\tilde{e}_i$ for $i = n{-}1, n$. Besides, we can check that φ preserves all of the (r, s)-Serre relations $(B5)$.

Case (iii): We claim that this case is impossible. First, we assume that $\varphi(\omega_{n-1})=\tilde\omega_{n-1}$ and $\varphi(\omega_n)=\tilde\omega_n'^{-1}$. Then we have $\varphi(e_{n-1})=a_1\tilde e_{n-1}$, $\varphi(e_n)=a_2\tilde f_n\tilde\omega_n'^{-1}$, where $a_1,a_2\in\mathbb{K}^*$. Using φ to relation $\omega_{n-1}e_n=s^2e_n\omega_{n-1}$, we get $s^2=s'^{-2}$; to relation $\omega_ne_{n-1}=r^{-2}e_{n-1}\omega_n$ to get $r^{-2}=s'^2$. So we get $r^2=s^2$. This is a contradiction.

Next, we assume that $\varphi(\omega_{n-1})=\tilde\omega_n$, and $\varphi(\omega_n)=\tilde\omega_{n-1}'^{-1}$. Then we have $\varphi(e_{n-1})=a_1\tilde e_n$, $\varphi(e_n)=a_2\tilde f_{n-1}\tilde\omega_{n-1}'^{-1}$, where $a_1,a_2\in\mathbb{K}^*$. Using φ to relation $\omega_{n-1}e_n=s^2e_n\omega_{n-1}$, we get $s^2=r'^2$; to relation $\omega_ne_{n-1}=r^{-2}e_{n-1}\omega_n$ to get $r^{-2}=r'^{-2}$. So we have $r^2=s^2$, which is contrary to the assumption.

Case (iv): Similarly to case (iii), it is also impossible.

(II) It suffices to consider the restriction φ_i of φ to an $\mathfrak{u}_{r_i,s_i}(\mathfrak{sl}_3)$-copy in $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ generated by $e_j,f_j,\omega_j^{\pm1},\omega_j'^{\pm1}$ for $j\in\{i,i{+}1\}$, where $1\le i<n{-}1$. Note that $\varphi(\omega_j)\in\{\tilde\omega_i,\tilde\omega_{i+1},\tilde\omega_i'^{-1},\tilde\omega_{i+1}'^{-1}\}$ for $j=i,i{+}1$.

Similarly to the proof of case (iii) or (iv) in (I), there don't exist the possibilities: $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_{j_i},\tilde\omega_{j_{i+1}}'^{-1})$, and $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_{j_i}'^{-1},\tilde\omega_{j_{i+1}})$ for $j_i\ne j_{i+1}\in\{i,i{+}1\}$. Thereby, we only need to consider the following possibilities: case (1): $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_i,\tilde\omega_{i+1})$; case (2): $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_i'^{-1},\tilde\omega_{i+1}'^{-1})$; case (3): $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_{i+1},\tilde\omega_i)$; case (4): $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_{i+1}'^{-1},\tilde\omega_i'^{-1})$.

It is easy to treat the first two cases:

Case (1): As $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_i,\tilde\omega_{i+1})$, by the argument of the paragraph above (I), we have $(\varphi(e_i),\varphi(e_{i+1}))=(a_i\tilde e_i,a_{i+1}\tilde e_{i+1})$ and $(\varphi(f_i),\varphi(f_{i+1}))=(a_i^{-1}\tilde f_i, a_{i+1}^{-1}\tilde f_{i+1})$, where $a_i,a_{i+1}\in\mathbb{K}^*$. By a similar argument of case (i) in (I), we have $(r'^2,s'^2)=(r^2,s^2)$.

Case (2): As $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_i'^{-1},\tilde\omega_{i+1}'^{-1})$, by the argument of the paragraph above (I), we have $(\varphi(e_i),\varphi(e_{i+1}))=(a_i\tilde f_i\tilde\omega_i'^{-1},a_{i+1}\tilde f_{i+1}\tilde\omega_{i+1}'^{-1})$ and $(\varphi(f_i),\varphi(f_{i+1}))$ $=(a_i^{-1}\tilde\omega_i^{-1}\tilde e_i,a_{i+1}^{-1}\tilde\omega_{i+1}^{-1}\tilde e_{i+1})$, where $a_i,a_{i+1}\in\mathbb{K}^*$. By a similar argument of case (ii) in (I), we have $(r'^2,s'^2)=(s^2,r^2)$.

As for the latter two cases, we have

Case (3): As $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_{i+1},\tilde\omega_i)$, using φ to $\omega_je_k=\langle\omega_k',\omega_j\rangle e_k\omega_j$ for $j,k\in\{i,i{+}1\}$, we get $\langle\tilde\omega_j',\tilde\omega_k\rangle=\langle\omega_k',\omega_j\rangle$. More precisely, we have $r'^2s'^{-2}=r^2s^{-2}$, $r'^{-2}=s^2$, $s'^2=r^{-2}$, i.e., $(r'^2,s'^2)=(s^{-2},r^{-2})$. Hence, $(\varphi(e_i),\varphi(e_{i+1}))=(a_{i+1}\tilde e_{i+1},a_i\tilde e_i)$ for $a_i,a_{i+1}\in\mathbb{K}^*$, $(\varphi(f_i),\varphi(f_{i+1}))=(rs)^{-2}(a_{i+1}^{-1}\tilde f_{i+1},a_i^{-1}\tilde f_i)$, since φ preserves $(B4)$.

Case (4): As $(\varphi(\omega_i),\varphi(\omega_{i+1}))=(\tilde\omega_{i+1}'^{-1},\tilde\omega_i'^{-1})$, using φ to $\omega_je_k=\langle\omega_k',\omega_j\rangle e_k\omega_j$ for $j,k\in\{i,i{+}1\}$, we get $\langle\tilde\omega_k',\tilde\omega_j\rangle^{-1}=\langle\omega_k',\omega_j\rangle$. More precisely, we have $r'^{-2}s'^2=r^2s^{-2}$, $r'^2=r^{-2}$, $s'^{-2}=s^2$, i.e., $(r'^2,s'^2)=(r^{-2},s^{-2})$. Hence, $(\varphi(e_i),\varphi(e_{i+1}))=(a_{i+1}\tilde f_{i+1}\tilde\omega_{i+1}'^{-1},a_i\tilde f_i\tilde\omega_i'^{-1})$ for $a_i,a_{i+1}\in\mathbb{K}^*$, $(\varphi(f_i),\varphi(f_{i+1}))=(rs)^{-2}(a_{i+1}^{-1}\tilde\omega_{i+1}^{-1}\tilde e_{i+1}, a_i^{-1}\tilde\omega_i^{-1}\tilde e_i)$, since φ preserves $(B4)$. Moreover, we can verify that φ preserves all of the (r,s)-Serre relations $(B5)$.

In view of the above discussions in cases (1)—(4), we derive the complete description below on all the Hopf algebra isomorphisms of $\mathfrak{u}_{r,s}(\mathfrak{sl}_n)$ (some of cases were missed in [**BW3**]).

Theorem 5.5. *Assume that rs^{-1} and $r's'^{-1}$ are primitive ℓth roots of unity. Then $\varphi:\ \mathfrak{u}_{r,s}(\mathfrak{sl}_n)\cong\mathfrak{u}_{r',s'}(\mathfrak{sl}_n)$ as Hopf algebras if and only if either*

(1) $(r', s') = (r, s)$, *φ is a diagonal isomorphism*: $\varphi(\omega_i) = \tilde{\omega}_i$, $\varphi(\omega'_i) = \tilde{\omega}'_i$, $\varphi(e_i) = a_i\tilde{e}_i$, $\varphi(f_i) = a_i^{-1}\tilde{f}_i$; *or*

(2) $(r', s') = (s, r)$, $\varphi(\omega_i) = \tilde{\omega}'^{-1}_i$, $\varphi(\omega'_i) = \tilde{\omega}_i^{-1}$, $\varphi(e_i) = a_i\tilde{f}_i\tilde{\omega}'^{-1}_i$, $\varphi(f_i) = a_i^{-1}\tilde{\omega}_i^{-1}\tilde{e}_i$; *or*

(3) $(r', s') = (s^{-1}, r^{-1})$, $\varphi(\omega_i) = \tilde{\omega}_{n-i}$, $\varphi(\omega'_i) = \tilde{\omega}'_{n-i}$, $\varphi(e_i) = a_{n-i}\tilde{e}_{n-i}$, $\varphi(f_i) = (rs)^{-1}a_{n-i}^{-1}\tilde{f}_{n-i}$; *or*

(4) $(r', s') = (r^{-1}, s^{-1})$, $\varphi(\omega_i) = \tilde{\omega}'^{-1}_{n-i}$, $\varphi(\omega'_i) = \tilde{\omega}^{-1}_{n-i}$, $\varphi(e_i) = a_{n-i}\tilde{f}_{n-i}\tilde{\omega}'^{-1}_{n-i}$, $\varphi(f_i) = (rs)^{-1}a_{n-i}^{-1}\tilde{\omega}^{-1}_{n-i}\tilde{e}_{n-i}$, $(a_i \in \mathbb{K}^*)$.

Now continue to the proof of Theorem 5.4. According to the discussions in (I), there are 2 possibilities: (i) $\varphi(\omega_{n-1}) = \tilde{\omega}_{n-1}$; (ii) $\varphi(\omega_{n-1}) = \tilde{\omega}'^{-1}_{n-1}$. Combining with the discussions in (II), we obtain two kinds of the Hopf algebra isomorphisms of $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ as required, which are only compatible with the cases (1) and (2) in (II), respectively. □

6. $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ is a Drinfel'd double

6.1. Let θ be a primitive ℓth root of unity in $\mathbb{K}$, and write $r = \theta^y$, $s = \theta^z$.

LEMMA 6.1. *Assume that $(2^{n-1}(y^n + (-1)^n z^n), \ell) = 1$ and rs^{-1} is a primitive ℓth root of unity with $\ell \neq 3$. Then $(\mathfrak{b}')^{coop} \cong \mathfrak{b}^*$ as Hopf algebras.*

PROOF. Define $\gamma_j \in \mathfrak{b}^*$ such that γ_j's are algebra homomorphisms with

$$\gamma_j(e_i) = 0, \ \forall\, i, \ \text{ and } \ \gamma_j(\omega_i) = \langle \omega'_j, \omega_i\rangle.$$

So they are group-like elements in $\mathfrak{b}^*$. Define $\eta_j = \sum\limits_{g \in G(\mathfrak{b})} (e_j g)^*$, where $G(\mathfrak{b})$ is the group generated by ω_i $(1 \leq i \leq n)$ and the asterisk denotes the dual basis element relative to the PBW-basis of $\mathfrak{b}$. The isomorphism $\phi : (\mathfrak{b}')^{coop} \to \mathfrak{b}^*$ is defined by

$$\phi(\omega'_j) = \gamma_j, \qquad \phi(f_j) = \eta_j.$$

First, we will check that ϕ is a Hopf algebra homomorphism and then we will show that it is a bijection.

Clearly, the γ_j's are invertible elements in $\mathfrak{b}^*$ that commute with one another and $\gamma_j^\ell = 1$. Note that $\eta_j^\ell = 0$, as it is 0 on any basis element of $\mathfrak{b}$. We calculate $\gamma_j\eta_i\gamma_j^{-1}$: It is nonzero only on basis elements of the form $e_i\omega_1^{k_1}\cdots\omega_n^{k_n}$, and on such an element it takes the value

$$\begin{aligned}
&(\gamma_j \otimes \eta_i \otimes \gamma_j^{-1})((e_i \otimes 1 \otimes 1 + \omega_i \otimes e_i \otimes 1 + \omega_i \otimes \omega_i \otimes e_i)(\omega_1^{k_1}\cdots\omega_n^{k_n})^{\otimes 3})\\
&= \gamma_j(\omega_i\omega_1^{k_1}\cdots\omega_n^{k_n})\eta_i(e_i\omega_1^{k_1}\cdots\omega_n^{k_n})\gamma_j^{-1}(\omega_1^{k_1}\cdots\omega_n^{k_n})\\
&= \gamma_j(\omega_i).
\end{aligned}$$

So, we have $\gamma_j\eta_i\gamma_j^{-1} = \langle\omega'_j, \omega_i\rangle\,\eta_i$, which corresponds to relation $(B3)$ for $\mathfrak{b}'$.

Next, with the same argument in the proof of Lemma 5.1 in [**HW**] (see, p. 263–264), we easily check relations in $(B5)$:

$$r^2s^2\eta_i^2\eta_{i+1} - (r^2{+}s^2)\eta_i\eta_{i+1}\eta_i + \eta_{i+1}\eta_i^2 = 0.$$

$$\eta_i\eta_{i+1}^2 - (r^{-2}{+}s^{-2})\eta_{i+1}\eta_i\eta_{i+1} + (rs)^{-2}\eta_{i+1}^2\eta_i = 0, \qquad i < n{-}1.$$

$$\begin{aligned}
&\eta_{n-1}\eta_n^3 - (r_n^{-2}{+}r_n^{-1}s_n^{-1}{+}s_n^{-2})\,\eta_n\eta_{n-1}\eta_n^2\\
&+ (r_ns_n)^{-1}(r_n^{-2}{+}r_n^{-1}s_n^{-1}{+}s_n^{-2})\,\eta_n^2\eta_{n-1}\eta_n - (r_ns_n)^{-3}\,\eta_n^3\eta_{n-1} = 0.
\end{aligned}$$

Hence, ϕ is an algebra homomorphism.

We have already seen that γ_i is a group-like element in $\mathfrak{b}^*$. Now using the same argument in the proof of Lemma 5.1 in [**HW**] (see, p. 264–265), we have

$$\Delta(\eta_i) = \eta_i \otimes 1 + \gamma_i \otimes \eta_i,$$

which means that ϕ is a Hopf algebra homomorphism.

Finally, we prove ϕ is bijective. As dim $\mathfrak{b}^* =$ dim $\mathfrak{b}'^{coop}$, it suffices to show ϕ is injective. By [**M**], we need only to show $\phi|_{(\mathfrak{b}')_1^{coop}}$ is injective. Lemma 5.2 yields $(\mathfrak{b}')_1^{coop} = \mathbb{K}G(\mathfrak{b}') + \sum_{i=1}^{n} \mathbb{K} f_i G(\mathfrak{b}')$, where $G(\mathfrak{b}')$ is the group generated by ω_i' $(1 \le i \le n)$. First, we claim that

$$\text{span}_{\mathbb{K}}\Big\{ \gamma_1^{k_1} \cdots \gamma_n^{k_n} \;\Big|\; 0 \le k_i < \ell \Big\} = \text{span}_{\mathbb{K}}\Big\{ (\omega_1^{k_1} \cdots \omega_n^{k_n})^* \;\Big|\; 0 \le k_i < \ell \Big\}. \tag{6.1}$$

This is equivalent to saying that the $\gamma_1^{k_1} \cdots \gamma_n^{k_n}$'s span the space of characters over $\mathbb{K}$ of the finite group $\mathbb{Z}/\ell\mathbb{Z} \times \cdots \times \mathbb{Z}/\ell\mathbb{Z}$ generated by $\omega_1, \cdots, \omega_n$. Since we assumed that $\mathbb{K}$ contains a primitive ℓth root of unity, the irreducible characters of this group are the functions $\chi_{i_1,\cdots,i_n}$ given by

$$\chi_{i_1,\cdots,i_n}(\omega_1^{k_1} \cdots \omega_n^{k_n}) = \theta^{i_1 k_1 + \cdots + i_n k_n},$$

where θ is a primitive ℓth root of unity in $\mathbb{K}$. Note that

$$\begin{aligned} \gamma_1 &= \chi_{2(y-z),-2y,0,\cdots,0}, \\ \gamma_2 &= \chi_{2z,2(y-z),-2y,0,\cdots,0}, \\ &\vdots \\ \gamma_{n-1} &= \chi_{0,\cdots,0,2z,2(y-z),-2y}, \\ \gamma_n &= \chi_{0,\cdots,0,2z,y-z}. \end{aligned}$$

We must show that, given $i_1, \cdots, i_n$, there are $k_1, \cdots, k_n$ such that

$$\chi_{i_1,\cdots,i_n} = \gamma_1^{k_1} \cdots \gamma_n^{k_n},$$

which is equivalent to the existence of a solution to the matrix equation

$$AK = I$$

in $\mathbb{Z}/\ell\mathbb{Z}$ (as these are powers of θ), where

$$A = \begin{pmatrix} 2(y-z) & 2z & 0 & 0 & \cdots & 0 & 0 \\ -2y & 2(y-z) & 2z & 0 & \cdots & 0 & 0 \\ 0 & -2y & 2(y-z) & 2z & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 2z & 0 \\ 0 & 0 & 0 & 0 & \cdots & 2(y-z) & 2z \\ 0 & 0 & 0 & 0 & \cdots & -2y & y-z \end{pmatrix},$$

K is the transpose of $(k_1, \cdots, k_n)$ and I is the transpose of $(i_1, \cdots, i_n)$. The determinant of the coefficient matrix A is $2^{n-1}(y^n + (-1)^n z^n)$, which is invertible in $\mathbb{Z}/\ell\mathbb{Z}$ by the hypothesis in the Lemma. So, (6.1) holds. In particular, this implies that the matrix

$$\big((\gamma_1^{k_1} \cdots \gamma_n^{k_n})(\omega_1^{j_1} \cdots \omega_n^{j_n}) \big)_{\bar{k} \times \bar{j}} \tag{6.2}$$

is invertible, and that ϕ is bijection on group-like elements.

Next, we will show for each i $(1 \leq i \leq n)$ that the following matrix is invertible:

$$\big((\eta_i \gamma_1^{k_1} \cdots \gamma_n^{k_n})(e_i \omega_1^{j_1} \cdots \omega_n^{j_n})\big)_{\bar{k} \times \bar{j}}\,. \tag{6.3}$$

This will complete the proof that ϕ is injective on $(\mathfrak{b}')_1^{coop}$, as desired. We will show that the matrix is block upper-triangular. Each matrix entry is

$$\begin{aligned}&(\eta_i \otimes \gamma_1^{k_1} \cdots \gamma_n^{k_n})(\Delta(e_i)\Delta(\omega_1^{j_1} \cdots \omega_n^{j_n}))\\ &\quad = (\eta_i \otimes \gamma_1^{k_1} \cdots \gamma_n^{k_n})(e_i \omega_1^{j_1} \cdots \omega_n^{j_n} \otimes \omega_1^{j_1} \cdots \omega_n^{j_n}).\end{aligned}$$

Thus, (6.3) is precisely the invertible matrix (6.2). □

6.2. With the same argument of Theorem 4.8 in [**BW3**], we have

Theorem 6.2. *Assume* $(2^{n-1}(y^n + (-1)^n z^n), \ell) = 1$, *then* $D(\mathfrak{b}) \cong \mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ *as Hopf algebras.* □

7. Integrals

7.1. Recall some definitions. Let H be a finite-dimensional Hopf algebra. An element $y \in H$ is a *left* (resp., *right*) *integral* if $ay = \varepsilon(a)y$ (resp., $ya = \varepsilon(a)y$) for all $a \in H$. The left (resp., right) integrals form a one-dimensional ideal $\int_H^l$ (resp., $\int_H^r$) of H, and $S_H(\int_H^r) = \int_H^l$ under the antipode S_H of H.

When $y \neq 0$ is a left integral of H, there exists a unique group-like element γ in the dual algebra H^* (the so-called *distinguished group-like element* of H^*) such that $ya = \gamma(a)y$. If we had begun instead with a right integral $y' \in H$, then we would have $ay' = \gamma^{-1}(a)y'$. This follows from the fact that group-like elements are invertible, and the following calculation, which can be found in [**M**, p.22]: When $y' \in \int_H^r$, $S_H(y')$ is a nonzero multiple of y, so that $S_H(y')S_H(a) = \gamma(S_H(a))S_H(y')$ for all $a \in H$. Applying S_H^{-1}, we find $ay' = \gamma(S_H(a))y'$. As γ is group-like, we have $\gamma(S_H(a)) = S_{H^*}(\gamma)(a) = \gamma^{-1}(a)$.

Now if $\lambda \neq 0$ is right integral of H^*, then there exists a unique group-like element g of H (the *distinguished group-like element* of H) such that $\xi\lambda = \xi(g)\lambda$ for all $\xi \in H^*$. The algebra H is *unimodular* (i.e., $\int_H^l = \int_H^r$) if and only if $\gamma = \varepsilon$; and the dual algebra H^* is unimodular if and only if $g = 1$.

The left and right H^*-module actions on H are given by

$$\xi \rightharpoonup a = \sum a_{(1)}\xi(a_{(2)}), \qquad a \leftharpoonup \xi = \sum \xi(a_{(1)})a_{(2)}, \tag{7.1}$$

for all $\xi \in H^*$ and $a \in H$. In particular, $\varepsilon \rightharpoonup a = a = a \leftharpoonup \varepsilon$ for all $a \in H$.

7.2. The first aim of this section is to construct the left and right integrals in the Borel subalgebra $\mathfrak{b}$ of $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$. To this end, we introduce the elements:

$$t = \prod_{i=1}^{n}(1 + \omega_i + \cdots + \omega_i^{\ell-1}), \quad x = \prod_{i=1}^{n} \mathcal{E}_{\widehat{i,i+1}},$$

where $\mathcal{E}_{\widehat{i,i+1}} = \mathcal{E}_{i,i}^{\ell-1}\mathcal{E}_{i,i+1}^{\ell-1} \cdots \mathcal{E}_{i,n}^{\ell-1}\mathcal{E}_{i,n'}^{\ell-1} \cdots \mathcal{E}_{i,i+1'}^{\ell-1}$ $(1 \leq i < n)$, $\mathcal{E}_{\widehat{n,n+1}} = \mathcal{E}_{n,n}^{\ell-1}$.

Theorem 7.1. *The element* $y = tx$ *is a left integral in* $\mathfrak{b}$.

Proof. We need to show that $by = \varepsilon(b)y$ for all $b \in \mathfrak{b}$. It suffices to show this for the generators ω_k and e_k, as the counit ε is an algebra homomorphism.

Observe that $\omega_k t = t = \varepsilon(\omega_k)t$ for all $k = 1, \cdots, n$, as the ω_i's commute and $\omega_k(1+\omega_k+\cdots+\omega_k^{\ell-1}) = 1+\omega_k+\cdots+\omega_k^{\ell-1}$. From that, relation $\omega_k y = \varepsilon(\omega_k)y$ is clear, for all k.

Next we compute $e_k y$. By a direct calculation, we get

$$\begin{aligned} e_k t = \prod_{i=1}^{n-1} \big(1 + r^{-2(\epsilon_i,\alpha_k)} s^{-2(\epsilon_{i+1},\alpha_k)} \omega_i + \cdots + r^{-2(\ell-1)(\epsilon_i,\alpha_k)} s^{-2(\ell-1)(\epsilon_{i+1},\alpha_k)} \omega_i^{\ell-1}\big) \cdot \\ \cdot \Big[\delta_{kn}(1 + r^{-1}s\omega_n + \cdots + r^{-(\ell-1)} s^{\ell-1} \omega_n^{\ell-1}) \\ + (1-\delta_{kn})(1 + r^{-2(\epsilon_n,\alpha_k)}\omega_n + \cdots + r^{-2(\ell-1)(\epsilon_n,\alpha_k)}\omega_n^{\ell-1})\Big] e_k. \end{aligned}$$

So it suffices to show that $e_k x = 0 = \varepsilon(e_k)x$.

Now

$$x = \mathcal{E}_{\widehat{1,2}}\mathcal{E}_{\widehat{2,3}} \cdots \mathcal{E}_{\widehat{k-1,k}}(\mathcal{E}_{k,k}^{\ell-1} \cdots \mathcal{E}_{k,k+1'}^{\ell-1}) \cdots \mathcal{E}_{\widehat{n,n+1}},$$
$$e_1 x = 0.$$

We want to show $e_k x = 0$, that is, e_k $(2 \le k \le n)$ can be moved across the terms $\mathcal{E}_{\widehat{i,i+1}}$ $(1 \le i < k)$ next to $\mathcal{E}_{k,k}^{\ell-1}$. By Lemma 3.1 (1), e_k commutes with $\mathcal{E}_{i,i}^{\ell-1} \cdots \mathcal{E}_{i,k-2}^{\ell-1}$ when $i \le k-2 \le n-2$. By Lemmas 3.9 (2) & 3.3 (5), we obtain

$$e_k \mathcal{E}_{i,k-1}^{\ell-1} = r^2 \mathcal{E}_{i,k-1}^{\ell-1} e_k + s^2 \mathcal{E}_{i,k-1}^{\ell-2} \mathcal{E}_{i,k}, \qquad (k \le n).$$

The second term reaches $\mathcal{E}_{i,k}^{\ell-1}$ and we get $\mathcal{E}_{i,k}^{\ell} = 0$. Hence, it suffices to treat the first term $r^2 \mathcal{E}_{i,k-1}^{\ell-1} e_k$ $(k \le n)$.

(i) For $k < n$: by Lemmas 3.2 (3), 3.3 (1) & (6), e_k quasi-commutes with $\mathcal{E}_{i,k}^{\ell-1} \mathcal{E}_{i,k+1}^{\ell-1} \cdots \mathcal{E}_{i,k+2'}^{\ell-1}$ (up to a factor s^2). By Lemmas 3.10 (3) & 3.6 (5), we obtain

$$e_k \mathcal{E}_{i,k+1'}^{\ell-1} = s^{-2} \mathcal{E}_{i,k+1'}^{\ell-1} e_k + r^{-2} \mathcal{E}_{i,k+1'}^{\ell-2} \mathcal{E}_{i,k'}, \qquad (k \le n-1).$$

The second term meets $\mathcal{E}_{i,k'}^{\ell-1}$ and yields $\mathcal{E}_{i,k'}^{\ell} = 0$. By Lemma 3.2 (5), we have $e_k \mathcal{E}_{i,k'}^{\ell-1} = r^{-2} \mathcal{E}_{i,k'}^{\ell-1} e_k$ $(i < k < n)$. This implies that e_k quasi-commutes with $\mathcal{E}_{\widehat{i,i+1}}$ for all $i < k$. Since $e_k \mathcal{E}_{\widehat{k,k+1}} = 0$, $e_k x = 0$ for $k < n$.

(ii) For $k = n$: by Lemmas 3.9 (4) & 3.5 (3), we have $e_n \mathcal{E}_{i,n}^{\ell-1} = (rs)\mathcal{E}_{i,n}^{\ell-1} e_n + s^2 \mathcal{E}_{i,n}^{\ell-2} \mathcal{E}_{i,n'}$. The second term meets $\mathcal{E}_{i,n'}^{\ell-1}$ and yields $\mathcal{E}_{i,n'}^{\ell} = 0$, so it remains to observe that e_n quasi-commutes with $\mathcal{E}_{i,n'}^{\ell-1} \mathcal{E}_{i,n-1'}^{\ell-1} \cdots \mathcal{E}_{i,i+1'}^{\ell-1}$ (by Lemmas 3.2 (4) & 3.4 (4)), that is, e_n quasi-commutes with those $\mathcal{E}_{\widehat{i,i+1}}$ for $i < n$ and finally meets $\mathcal{E}_{\widehat{n,n+1}}$ and yields 0.

This completes the proof. □

Theorem 7.2. *The element $y' = xt$ is a right integral in $\mathfrak{b}$.*

Proof. Arguing as in Theorem 7.1, we see that $t\omega_j = t = \varepsilon(\omega_j)t$, and hence that $y'\omega_j = \varepsilon(\omega_j)y'$ for all j. As

$$\begin{aligned} te_j = e_j \prod_{i=1}^{n-1}(1 + r^{2(\epsilon_i,\alpha_j)}s^{2(\epsilon_{i+1},\alpha_j)}\omega_i + \cdots + r^{2(\ell-1)(\epsilon_i,\alpha_j)}s^{2(\ell-1)(\epsilon_{i+1},\alpha_j)}\omega_i^{\ell-1})\cdot \\ \cdot\Big[\delta_{jn}e_j(1 + rs^{-1}\omega_n + \cdots + r^{\ell-1}s^{-(\ell-1)}\omega_n^{\ell-1}) \\ + (1-\delta_{jn})\, e_j(1 + r^{2(\epsilon_n,\alpha_j)}\omega_n + \cdots + r^{2(\ell-1)(\epsilon_n,\alpha_j)}\omega_n^{\ell-1})\Big], \end{aligned}$$

so it suffices to show that $xe_j = 0$.

Now

$$x = \mathcal{E}_{\widehat{1,2}} \cdots \mathcal{E}_{\widehat{j,j+1}}\Big(\mathcal{E}^{\ell-1}_{j+1,j+1}\cdots\mathcal{E}^{\ell-1}_{j+1,k'}\cdots\mathcal{E}^{\ell-1}_{j+1,j+2'}\Big)\cdots\mathcal{E}_{\widehat{n,n+1}},$$

$$xe_n = 0.$$

We need to prove $xe_j = 0$ for $1 \le j < n$. By Lemmas 3.9 (3), (4), (6) & 3.4 (3), it is easy to see that $\mathcal{E}_{\widehat{n-1,n}}\mathcal{E}_{\widehat{n,n+1}}e_{n-1} = 0$. It suffices to show $\mathcal{E}_{\widehat{j,j+1}}\mathcal{E}_{\widehat{j+1,j+2}}e_j = 0$ since e_j commutes with $\mathcal{E}_{\widehat{j+2,j+3}}\cdots\mathcal{E}_{\widehat{n,n+1}}$ (by Lemma 3.1 (1)) for $j < n-1$.

(I) Claim: $\mathcal{E}_{\widehat{j,j+1}}e_j = 0$ for $j < n{-}1$.

Write

$$\begin{aligned} \Pi_j^{(k)} &:= \mathcal{E}^{\ell-1}_{j,j}\mathcal{E}^{\ell-1}_{j,j+1}\cdots\mathcal{E}^{\ell-1}_{j,k}, \quad (j \le k \le n), \\ \Pi_j^{(k')} &:= \Pi_j^{(n)}\,\mathcal{E}^{\ell-1}_{j,n'}\cdots\mathcal{E}^{\ell-1}_{j,k'}, \quad (n' \le k' \le j{+}1'). \end{aligned}$$

Thus, $\Pi_j^{(j+1')} = \mathcal{E}_{\widehat{j,j+1}}$, and $\Pi_j^{(j+2')}e_j = 0$, as $e_j\mathcal{E}_{j,k} = s^2\mathcal{E}_{j,k}e_j$ for $j < k \le n$, $e_j\mathcal{E}_{j,k'} = s^2\mathcal{E}_{j,k'}e_j$ for $j{+}1 < k \le n$, and $e_j^\ell = 0$.

For $j < n{-}1$, by Lemma 3.10 (4) & (5) and $\mathcal{E}^\ell_{j,j+2'} = \cdots = \mathcal{E}^\ell_{j,n'} = \mathcal{E}^\ell_{j,n} = 0$, we get

$$\begin{aligned} \Pi_j^{(j+1')}e_j &= \Pi_j^{(j+2')}\Big(\mathcal{E}^{\ell-1}_{j,j+1'}e_j\Big) && \text{(by Lemma 3.10 (4))} \\ &= *_1\Pi_j^{(j+2')}\mathcal{E}^{\ell-1}_{j+1,j+1}\cdots\mathcal{E}^{\ell-1}_{j+1,n-2}\cdots\,\mathcal{E}_{j,j+1}\mathcal{E}_{j,j+2'}\mathcal{E}^{\ell-2}_{j,j+1'} \\ &= *_2\Pi_j^{(j+3')}\Big(\mathcal{E}^{\ell-1}_{j,j+2'}\mathcal{E}_{j,j+1}\Big)\mathcal{E}_{j,j+2'}\mathcal{E}^{\ell-2}_{j,j+1'} && \text{(by Lemma 3.10 (4))} \\ &= *_2\Pi_j^{(j+3')}\Big(\mathcal{E}_{j,j+1}\mathcal{E}^{\ell-1}_{j,j+2'} - (rs)^{-2}\mathcal{E}_{j,j+2}\mathcal{E}_{j,j+3'}\mathcal{E}^{\ell-2}_{j,j+2'}\Big)\mathcal{E}_{j,j+2'}\mathcal{E}^{\ell-2}_{j,j+1'} \\ &= *_3\Pi_j^{(j+4')}\Big(\mathcal{E}^{\ell-1}_{j,j+3'}\mathcal{E}_{j,j+2}\Big)\mathcal{E}_{j,j+3'}\mathcal{E}^{\ell-1}_{j,j+2'}\mathcal{E}^{\ell-2}_{j,j+1'} \\ &= \cdots && \text{(by Lemma 3.10 (4))} \\ &= *\,\Pi_j^{(n)}\Big(\mathcal{E}^{\ell-1}_{j,n'}\mathcal{E}_{j,n-1}\Big)\mathcal{E}_{j,n'}\mathcal{E}^{\ell-1}_{j,n-1'}\cdots\mathcal{E}^{\ell-1}_{j,j+2'}\mathcal{E}^{\ell-2}_{j,j+1'} && \text{(by Lemma 3.10 (5))} \\ &= *'\Pi_j^{(n)}\Big(\mathcal{E}^2_{j,n}\mathcal{E}^{\ell-2}_{j,n'}\Big)\mathcal{E}_{j,n'}\mathcal{E}^{\ell-1}_{j,n-1'}\cdots\mathcal{E}^{\ell-1}_{j,j+2'}\mathcal{E}^{\ell-2}_{j,j+1'} \\ &= 0, \qquad (\text{since } \Pi_j^{(n)}\mathcal{E}_{j,n} = 0). \end{aligned}$$

(II) Write $\Delta_j^{(k)} := \Pi_j^{(j+1')}\Pi_{j+1}^{(k)}$ for $j{+}1 \le k \le n$. Then $\Delta_j^{(n)} = \Pi_j^{(j+1')}\Pi_{j+1}^{(n)}$. We make a convention: $\Delta_j^{(j)} := \Pi_j^{(j+1')}$.

(i) Assume $\Delta_j^{(k)}e_j = 0$ for $j \le k < n$, we want to prove $\Delta_j^{(k+1)}e_j = 0$.

(1) For $k+1<n$: indeed, by Lemmas 3.9 (1) & 3.3 (4), we have

$$\begin{aligned}\Delta_j^{(k+1)}e_j &= \Delta_j^{(k)}\Big(\mathcal{E}_{j+1,k+1}^{\ell-1}e_j\Big)\\ &= \Delta_j^{(k)}\Big(r^2e_j\mathcal{E}_{j+1,k+1}+s^2\mathcal{E}_{j,k+1}\Big)\mathcal{E}_{j+1,k+1}^{\ell-2} \quad \text{(by Lemma 3.3 (3))}\\ &= s^2\Pi_j^{(j+1')}\mathcal{E}_{j,k+1}\Pi_{j+1}^{(k)}\mathcal{E}_{j+1,k+1}^{\ell-2}.\end{aligned}$$

Lemmas 3.7 (8), 3.5 (3), 3.3 (9) show that $\mathcal{E}_{j,k+1}$ quasi-commutes with any element of $\{\,\mathcal{E}_{j,l'}\mid j+1\le l\le n,\ l\ne k+2\,\}$. Hence, by a similar argument in (I) (using Lemma 3.10 (4) & (5)), we can get

$$\begin{aligned}\Pi_j^{(j+1')}\mathcal{E}_{j,k+1} &= *_1\,\Pi_j^{(k+3')}\Big(\mathcal{E}_{j,k+2'}^{\ell-1}\mathcal{E}_{j,k+1}\Big)\mathcal{E}_{j,k+1'}^{\ell-1}\cdots\mathcal{E}_{j,j+1'}^{\ell-1}\\ &= *_2\,\Pi_j^{(k+3')}\Big(\mathcal{E}_{j,k+1}\mathcal{E}_{j,k+2'}^{\ell-1}-(rs)^{-2}\mathcal{E}_{j,k+2}\mathcal{E}_{j,k+3'}\mathcal{E}_{j,k+2'}^{\ell-2}\Big)\cdots\mathcal{E}_{j,j+1'}^{\ell-1}\\ &= 0+*_3\,\Pi_j^{(k+4')}\Big(\mathcal{E}_{j,k+3'}^{\ell-1}\mathcal{E}_{j,k+2}\Big)\mathcal{E}_{j,k+3'}\mathcal{E}_{j,k+2'}^{\ell-2}\mathcal{E}_{j,k+1'}^{\ell-1}\cdots\mathcal{E}_{j,j+1'}^{\ell-1}\\ &= 0,\end{aligned}$$

here in the 1st summand 0 we used the fact that $\mathcal{E}_{j,k+1}$ quasi-commutes with $\mathcal{E}_{j,n'}^{\ell-1}\cdots\mathcal{E}_{j,k+3'}^{\ell-1}$, as well as $\mathcal{E}_{j,l}$ (by Lemma 3.3 (5)) for any $k+2\le l\le n$, and $\mathcal{E}_{j,k+1}^{\ell}=0$; while the 2nd summand 0 used repeatedly the proof-procedure of (I).

(2) For $k+1=n$: at first we note that

$$\Delta_j^{(n-1)}\mathcal{E}_{j,n}=0, \qquad \Delta_j^{(n-1)}\mathcal{E}_{j+1,n-1}=0,$$

since Lemmas 3.3 (3) & 3.5 (3) say: $\mathcal{E}_{j,n}$ commutes with $\mathcal{E}_{j+1,l}$ for $j+1\le l\le n-1$, and quasi-commutes with $\mathcal{E}_{j,l'}$ for $j+1\le l\le n$. Thus, by Lemma 3.4 (1), we have

$$\mathcal{E}_{j+1,n}\mathcal{E}_{j,n}\equiv rs^{-1}\mathcal{E}_{j,n'}\mathcal{E}_{j+1,n-1} \mod(\mathcal{E}_{j,n}\mathcal{E}_{j+1,n}+\mathcal{E}_{j+1,n-1}\mathcal{E}_{j,n'}). \tag{7.2}$$

Furthermore, we note that

$$\Delta_j^{(k-2)}\mathcal{E}_{j,k'}=0, \qquad \Delta_j^{(k-2)}\mathcal{E}_{j+1,k-2}=0, \qquad (j{+}2<k\le n). \tag{7.3}$$

Hence, by Lemma 3.4 (2), for $j{+}2<k\le n$, we get

$$\mathcal{E}_{j+1,k-1}\mathcal{E}_{j,k'}\equiv(rs)^2\mathcal{E}_{j,k-1'}\mathcal{E}_{j+1,k-2} \mod(\mathcal{E}_{j,k'}\mathcal{E}_{j+1,k-1}-\mathcal{E}_{j+1,k-2}\mathcal{E}_{j,k-1'}). \tag{7.4}$$

Using (7.2), (7.3) & (7.4), by Lemmas 3.9 (5), 3.3 (5) & 3.10 (6), we get

$$\begin{aligned}\Delta_j^{(n)}e_j = \Delta_j^{(n-1)}\big(\mathcal{E}_{j+1,n}^{\ell-1}e_j\big) &= \Delta_j^{(n-1)}\Big[r^2e_j\mathcal{E}_{j+1,n}^{\ell-1}+(rs)^{-1}\mathcal{E}_{j+1,n}^{\ell-2}\mathcal{E}_{j,n}\\ &\quad+(rs^3)^{-1}\mathcal{E}_{j+1,n}^{\ell-3}\Big(r^2\mathcal{E}_{j,n'}\mathcal{E}_{j+1,n-1}-\mathcal{E}_{j+1,n-1}\mathcal{E}_{j,n'}\Big)\Big] \quad \text{(by Lemma 3.3 (5))}\\ &= (rs)^{-1}\Delta_j^{(n-1)}\Big[\mathcal{E}_{j+1,n}^{\ell-2}\mathcal{E}_{j,n}+s^4\mathcal{E}_{j,n'}\mathcal{E}_{j+1,n-1}\mathcal{E}_{j+1,n}^{\ell-3}\Big] \quad \text{(by Lemma 3.5 (1))}\end{aligned}$$

$$
\begin{aligned}
&= \Delta_j^{(n-1)}\mathcal{E}_{j,n'}\mathcal{E}_{j+1,n-1}\Big(*_1\mathcal{E}_{j+1,n-1}^{\ell-3} + *_2\mathcal{E}_{j+1,n}^{\ell-3}\Big)\\
&= \Delta_j^{(n-2)}\Big(\mathcal{E}_{j+1,n-1}^{\ell-1}\mathcal{E}_{j,n'}\Big)\mathcal{E}_{j+1,n-1}(\cdots)\\
&= *_3\Delta_j^{(n-3)}\Big(\mathcal{E}_{j+1,n-2}^{\ell-1}\mathcal{E}_{j,n-1'}\Big)\mathcal{E}_{j+1,n-2}^{\ell-1}\mathcal{E}_{j+1,n-1}(\cdots)\\
&= \cdots \qquad\qquad\qquad\qquad \text{(using (7.3) \& (7.4) repeatedly)}\\
&= *\Delta_j^{(j)}\Big(\mathcal{E}_{j+1,j+1}^{\ell-1}\mathcal{E}_{j,j+2'}\Big)\Pi_{j+1}^{(n-2)}\mathcal{E}_{j+1,n-1}(\cdots) \qquad \text{(by Lemma 3.10 (6))}\\
&= *\Pi_j^{(j+1')}\Big(s^{-2}\mathcal{E}_{j,j+2'}e_{j+1} + r^{-2}\mathcal{E}_{j,j+1'}\Big)e_{j+1}^{\ell-2}\,\Pi_{j+1}^{(n-2)}\mathcal{E}_{j+1,n-1}(\cdots)\\
&= 0, \qquad \text{(since } e_{j+1}\Pi_{j+1}^{(n-2)} = 0).
\end{aligned}
$$

(ii) Write $\Delta_j^{(k')} := \Pi_j^{(j+1')}\Pi_{j+1}^{(k')}$ for $n' \le k' \le j+2'$, and make a convention: $\Delta_j^{(n+1')} := \Pi_j^{(j+1')}$. Assume $\Delta_j^{(k+1')}e_j = 0$ for $n+1' \le k+1' < j+2'$, we want to prove $\Delta_j^{(k')}e_j = 0$.

By Lemmas 3.10 (1) & 3.5 (6) or 3.6 (7), we obtain

$$
\begin{aligned}
&\mathcal{E}_{j+1,n'}^{\ell-1}e_j = r^2e_j\mathcal{E}_{j+1,n'}^{\ell-1} + (r^2s)^2\mathcal{E}_{j,n'}\mathcal{E}_{j+1,n'}^{\ell-2},\\
&\mathcal{E}_{j+1,k'}^{\ell-1}e_j = r^2e_j\mathcal{E}_{j+1,k'}^{\ell-1} + s^2\mathcal{E}_{j,k'}\mathcal{E}_{j+1,k'}^{\ell-2}, \qquad j+2 \le k < n.
\end{aligned}
\tag{7.5}
$$

By Lemmas 3.6 (6) & 3.5 (5), $\mathcal{E}_{j,k'}$ quasi-commutes with $\mathcal{E}_{j+1,n}^{\ell-1}\mathcal{E}_{j+1,n'}^{\ell-1}\cdots\mathcal{E}_{j+1,k+1'}^{\ell-1}$ (up to a factor of some power of $(rs)^2$). Again, Lemma 3.7 (7) indicates that $\mathcal{E}_{j,k'}$ quasi-commutes with $\mathcal{E}_{j+1,k}^{\ell-1}\cdots\mathcal{E}_{j+1,n-1}^{\ell-1}$ (up to a factor of some power of $(rs)^{-2}$). Based on these data, together with (7.3) & (7.4), using the same proof-procedure as the above (i): (2), we get

$$
\begin{aligned}
\Delta_j^{(k')}e_j &= \Delta_j^{(k+1')}\Big(\mathcal{E}_{j+1,k'}^{\ell-1}e_j\Big) && \text{(by (7.5))}\\
&= *\,\Delta_j^{(k-2)}\Big(\mathcal{E}_{j+1,k-1}^{\ell-1}\mathcal{E}_{j,k'}\Big)\mathcal{E}_{j+1,k}^{\ell-1}\cdots\mathcal{E}_{j+1,k'}^{\ell-1} && \text{(by (7.4))}\\
&= \cdots && \text{(using the same procedure of (i): (2))}\\
&= 0.
\end{aligned}
$$

The proof is complete. □

7.3. A finite-dimensional Hopf algebra H is semisimple if and only if $\varepsilon(\int_H^l) \neq 0$ or equivalently, $\varepsilon(\int_H^r) \neq 0$. For the algebra $\mathfrak{b}$ above, y gives a basis for $\int_{\mathfrak{b}}^l$ and y' a basis for $\int_{\mathfrak{b}}^r$. As $\varepsilon(y) = 0 = \varepsilon(y')$, we have

PROPOSITION 7.3. *The Hopf subalgebra* $\mathfrak{b}$ *is not semisimple.* □

7.4. The second aim of this section is to single out the distinguished group-like elements of $\mathfrak{b}$ and $\mathfrak{b}^*$. Since the group-like elements of $\mathfrak{b}^*$ are exactly the algebra homomorphisms in $\mathrm{Alg}_{\mathbb{K}}(\mathfrak{b}, \mathbb{K})$, it suffices to compute its values on the generators.

PROPOSITION 7.4. *Write* $2\rho = \sum_{j=1}^{n} j(2n-j)\alpha_j$, *where* ρ *is the half sum of positive roots of* $\mathfrak{so}_{2n+1}$. *Let* $\gamma \in Alg_{\mathbb{K}}(\mathfrak{b}, \mathbb{K})$ *be defined by*

$$
\gamma(e_k) = 0, \qquad \gamma(\omega_k) = \langle \omega'_{2\rho}, \omega_k\rangle.
\tag{7.6}
$$

Then γ *is the distinguished group-like element of* $\mathfrak{b}^*$.

PROOF. It suffices to argue that γ as in (7.6) satisfies $ya = \gamma(a)y$ for $a = e_k$ and $a = \omega_k$, $1 \leq k \leq n$, and for $y = tx$ given in Theorem 7.1. Recall from the proof of Theorem 7.2 that $xe_k = 0$. Thus, $ye_k = txe_k = 0 = \gamma(e_k)y$. We have

$$y\omega_k = t\Big(\prod_{i=1}^{n} \mathcal{E}_{\widehat{i,i+1}}\Big)\omega_k = \Pi_{i=1}^{n} \langle \omega_i', \omega_k \rangle^{i(2n-i)}\, t\omega_k x = \gamma(\omega_k)y.$$

This completes the proof. $\square$

Under the assumptions of Lemma 6.1, $(\mathfrak{b}')^{coop} \cong \mathfrak{b}^*$ as Hopf algebras, via the map $\phi : (\mathfrak{b}')^{coop} \to \mathfrak{b}^*, \phi(\omega_j') = \gamma_j, \phi(f_j) = \eta_j$ (for definition, see the proof of Lemma 6.1.). This allows us to define a Hopf pairing $\mathfrak{b}' \times \mathfrak{b} \to \mathbb{K}$ whose values on generators are given by

$$(f_j \,|\, e_i) = \delta_{ij}, \qquad (\omega_j' \,|\, \omega_i) = \langle \omega_j', \omega_i \rangle, \tag{7.7}$$

and are zero on all other pairs of generators.

By (7.6) & (7.7), we get $(\omega_{2\rho}' \,|\, b) = \gamma(b)$, for all $b \in \mathfrak{b}$.

Note that $\mathfrak{b}_{s^{-1},r^{-1}} \cong (\mathfrak{b}')^{coop} \cong \mathfrak{b}^*$ as Hopf algebras. Under the isomorphism $\phi\psi^{-1}$ (where $\psi(f_i) = e_i$, $\psi(\omega_i') = \omega_i$), a nonzero left (resp., right) integral of $\mathfrak{b}$ maps to a nonzero left (resp., right) integral of $\mathfrak{b}^*$. Thus, we have

PROPOSITION 7.5. *Let* $\lambda = \nu\eta$ *and* $\lambda' = \eta\nu \in \mathfrak{b}^*$, *where*

$$\nu = \prod_{i=1}^{n}(1 + \gamma_i + \cdots + \gamma_i^{\ell-1}), \quad \eta = \prod_{i=1}^{n} \eta_{\widehat{i,i+1}},$$

where $\eta_{\widehat{i,i+1}} = \eta_{i,i}^{\ell-1}\eta_{i,i+1}^{\ell-1}\cdots\eta_{i,n}^{\ell-1}\eta_{i,n'}^{\ell-1}\cdots\eta_{i,i+1'}^{\ell-1}$ $(i < n)$, $\eta_{\widehat{n,n+1}} = \eta_{n,n}^{\ell-1}$, $\eta_{i,i} = \eta_i$, $\eta_{i,j} = [\eta_{i+1,j}, \eta_i]_{s^2}$, $\eta_{i,n'} = [\eta_n, \eta_{i,n}]_{rs}$, $\eta_{i,j'} = [\eta_j, \eta_{i,j+1'}]_{r^{-2}}$.

Then λ *is a left integral and* λ' *is a right integral of* $\mathfrak{b}^*$. $\square$

COROLLARY 7.6. *The element* $g = \omega_{2\rho}^{-1}$ *is the distinguished group-like element of* $\mathfrak{b}$, *and under the Hopf pairing in* (7.7),

$$(\omega_i' \,|\, g) = \langle \omega_i', \omega_{2\rho}^{-1} \rangle = \gamma_i(g).$$

PROOF. Let $\mathcal{F} = \prod_{i=1}^{n} \mathcal{F}_{\widehat{i,i+1}}$, where $\mathcal{F}_{\widehat{i,i+1}} = \mathcal{F}_{i,i}^{\ell-1}\mathcal{F}_{i,i+1}^{\ell-1}\cdots\mathcal{F}_{i,n}^{\ell-1}\mathcal{F}_{i,n'}^{\ell-1}\cdots\mathcal{F}_{i,i+1'}^{\ell-1}$ for $i \leq n{-}1$, and $\mathcal{F}_{\widehat{n,n+1}} = \mathcal{F}_{n,n}^{\ell-1}$. Then we have

$$\omega_k'\mathcal{F} = \langle \omega_k', \omega_{2\rho} \rangle^{-1}\mathcal{F}\omega_k'.$$

Because $\phi^{-1}(\lambda') = \mathcal{F}\Big(\prod_{i=1}^{n}(1 + \omega_i' + \cdots + (\omega_i')^{\ell-1})\Big)$, it follows that

$$\gamma_k\lambda' = \langle \omega_k', \omega_{2\rho} \rangle^{-1}\lambda', \quad \text{and} \quad \eta_k\lambda' = 0.$$

Taking $g = \omega_{2\rho}^{-1}$, we have $\xi\lambda' = \xi(g)\lambda'$ for all $\xi \in \mathfrak{b}^*$. $\square$

8. To be a ribbon Hopf algebra

8.1. A finite-dimensional Hopf algebra H is *quasitriangular* if there is an invertible element $R = \sum x_i \otimes y_i$ in $H \otimes H$ such that $\Delta^{op}(a) = R\Delta(a)R^{-1}$ for all $a \in H$, and R satisfies the relations $(\Delta \otimes id)R = R_{1,3}R_{2,3}$, $(id \otimes \Delta)R = R_{1,3}R_{1,2}$, where $R_{1,2} = \sum x_i \otimes y_i \otimes 1$, $R_{1,3} = \sum x_i \otimes 1 \otimes y_i$, and $R_{2,3} = \sum 1 \otimes x_i \otimes y_i$.

Write $u = \sum S(y_i)x_i$. Then $c = uS(u)$ is central in H (cf. [**Ka**]) and is referred to as the *Casimir element*.

An element $v \in H$ is a *quasi-ribbon element* of quasitriangular Hopf algebra (H, R) if (i) $v^2 = c$, (ii) $S(v) = v$, (iii) $\varepsilon(v) = 1$, (iv) $\Delta(v) = (R_{2,1}R)^{-1}(v \otimes v)$, where $R_{2,1} = \sum y_i \otimes x_i$. If moreover v is central in H, then v is a *ribbon element*, and (H, R, v) is called a *ribbon Hopf algebra*. Ribbon elements provide a very effective means of constructing invariants of knots or links.

8.2. The Drinfel'd double $D(A)$ of a finite-dimensional Hopf algebra A is quasitriangular, and Kauffman-Radford ([**KR**]) proved a criterion for $D(A)$ to be ribbon.

Theorem 8.1. ([KR, Thm. 3]) *Assume A is a finite-dimensional Hopf algebra, let g and γ be the distinguished group-like elements of A and A^* respectively. Then*

(i) *$(D(A), R)$ has a quasi-ribbon element if and only if there exist group-like elements $h \in A$, $\delta \in A^*$ such that $h^2 = g$, $\delta^2 = \gamma$.*

(ii) *$(D(A), R)$ has a ribbon element if and only if there exist h and δ as in* (i) *such that $S^2(a) = h(\delta \rightharpoonup a \leftharpoonup \delta^{-1})h^{-1}$, $\forall\, a \in A$.*

8.3. As a consequence of Theorem 8.1, we have

Theorem 8.2. *Assume r, s are ℓth roots of unity. Then for the Hopf subalgebra $\mathfrak{b}$ of $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$, the following are equivalent*

(i) *$D(\mathfrak{b})$ has a quasi-ribbon element;*

(ii) *$D(\mathfrak{b})$ has a ribbon element;*

(iii) *ℓ is odd.*

Proof. By Corollary 7.6, $g = \omega_{2\rho}^{-1}$ is the distinguished group-like element of $\mathfrak{b}$. There exists a group-like element $h = \prod_{j=1}^{n} \omega_j^{a_j} \in \mathfrak{b}$ with $a_j \in \mathbb{Z}$ such that $h^2 = g$ if and only if the equations $2a_j \equiv -j(2n-j) \mod \ell$ can be solved for $j = 1, \cdots, n$. Solutions exist if and only if ℓ is odd. Suppose now ℓ is odd, then we can take $h = \omega_\rho^{-1} \in \mathfrak{b}$. Because $\gamma = (\omega'_{2\rho} \,|\, \cdot)$ corresponds to $\omega'_{2\rho}$ under the isomorphism $\phi^{-1} : \mathfrak{b}^* \to (\mathfrak{b}')^{coop}$, there exists a $\delta = (\omega'_\rho \,|\, \cdot) \in \mathfrak{b}^*$ such that $\delta^2 = \gamma$, where $\delta \in \mathrm{Alg}_{\mathbb{K}}(\mathfrak{b}, \mathbb{K})$ is defined by

$$\delta(e_k) = 0, \qquad \delta(\omega_k) = \langle \omega'_\rho, \omega_k \rangle.$$

Then using (7.1), we get

$$\begin{aligned} h(\delta \rightharpoonup \omega_k \leftharpoonup \delta^{-1})h^{-1} &= \delta(\omega_k)\delta^{-1}(\omega_k)h\omega_k h^{-1} = \omega_k = S^2(\omega_k). \\ h(\delta \rightharpoonup e_k \leftharpoonup \delta^{-1})h^{-1} &= \delta(1)\delta^{-1}(\omega_k)he_k h^{-1} \\ &= \langle \omega'_\rho, \omega_k \rangle^{-1} he_k h^{-1} \\ &= \langle \omega'_\rho, \omega_k \rangle^{-1} \langle \omega'_k, \omega_\rho \rangle^{-1} e_k \\ &= r_k^{-1} s_k e_k = \omega_k^{-1} e_k \omega_k = S^2(e_k). \end{aligned}$$

By Theorem 8.1, we complete the proof. □

By Theorem 6.2, we have

Corollary 8.3. *Assume that* $r = \theta^y$, $s = \theta^z$, *where* θ *is a primitive* ℓ*th root of unity and* $(2^{n-1}(y^n + (-1)^n z^n), \ell) = 1$. *Then the following are equivalent*

(i) $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ *has a quasi-ribbon element;*

(ii) $\mathfrak{u}_{r,s}(\mathfrak{so}_{2n+1})$ *has a ribbon element;*

(iii) ℓ *is odd.* □

References

[B] J. Beck, *Convex bases of PBW type for quantum affine algebras*, Comm. Math. Phys. **165** (1994), 193–199.

[BGH1] N. Bergeron, Y. Gao and N. Hu, *Drinfel'd doubles and Lusztig's symmetries of two-parameter quantum groups*, (arXiv:math/0505614), J. Algebra **301** (2006), 378–405.

[BGH2] N. Bergeron, Y. Gao and N. Hu, *Representations of two-parameter quantum orthogonal groups and symplectic groups*, (arXiv math. QA/0510124), AMS/IP, Studies in Advanced Mathematics, vol. **39**, pp. 1–21, 2007.

[BH] X. Bai and N. Hu, *Two-parameter quantum group of exceptional type E-series and convex PBW type basis*, (arXiv.math.QA/0605179), Algebra Colloq., **15** (4) (2008), 619–636.

[BW1] G. Benkart and S. Witherspoon, *Two-parameter quantum groups (of type A) and Drinfel'd doubles*, Algebra Represent. Theory, **7** (2004), 261–286.

[BW2] G. Benkart and S. Witherspoon, *Representations of two-parameter quantum groups (of type A) and Schur-Weyl duality*, in "Hopf Algebras", Lecture Notes in Pure and Appl. Math., **237**, pp. 65–92, Dekker, New York, 2004.

[BW3] G. Benkart and S. Witherspoon, *Restricted two-parameter quantum groups (of type A)*, Fields Institute Communications, "Representations of Finite Dimensional Algebras and Related Topics in Lie Theory and Geometry",vol. **40**, pp. 293–318, Amer. Math. Soc., Providence, RI, 2004.

[CX] V. Chari and N. Xi, *Monomial bases of quantized enveloping algebras*, in "Recent developments in quantum affine algebras and related topics", Contemp. Math. vol. **216**, pp. 23–57, Amer. Math. Soc., 2001.

[H] N. Hu, *Lyndon words, convex PBW bases and their R-matrices for the two-parameter quantum groups of* B_2, C_2, D_4 *types*, manuscript 2005.

[HP] N. Hu and Y. Pei, *Notes on two-parameter quantum groups, (I)*, (arXiv.math.QA/0702298), Sci. in China, Ser. A. **51** (6) (2008), 1101–1110.

[HRZ] N. Hu, M. Rosso and H. Zhang, *Two-parameter affine quantum group* $U_{r,s}(\widehat{\mathfrak{sl}_n})$, *Drinfel'd realization and quantum affine Lyndon basis*, (arXiv:0812.3107), Comm. Math. Phys. **278** (2) (2008), 453–486.

[HS] N. Hu and Q. Shi, *The two-parameter quantum group of exceptional type* G_2 *and Lusztig symmetries*, (arXiv:math/0601444), Pacific J. Math., **230** (2) (2007), 327–346.

[HW] N. Hu and X. Wang, *Convex PBW-type Lyndon basis and restricted two-parameter quantum groups of type* G_2, (arXiv:0811.0209), Pacific J. Math. **241** (2) (2009), 243–273.

[HZ] N. Hu and H. Zhang, *Vertex representations of two-parameter quantum affine algebras* $U_{r,s}(\widehat{\mathfrak{g}})$*: the simply-laced cases*, Preprint 2006-2007.

[Ka] C. Kassel, *Quantum Groups*, GTM **155**, Springer-Verlag Berlin/Heidelberg/New York, 1995.

[KR] L.H. Kauffman and D.E. Radford, *A necessary and sufficient condition for a finite-dimensional Drinfel'd double to be a ribbon Hopf algebra*, J. Algebra **159** (1993), 98–114.

[K] V.K. Kharchenko, *A combinatorial approach to the quantification of Lie algebras*, Pacific J. Math. **23** (2002), 191–233.

[LR] M. Lalonde and A. Ram, *Standard Lyndon bases of Lie algebras and enveloping algebras*, Trans. Amer. Math. Soc., **347** (5) (1995), 1821–1830.

[Lu] G. Lusztig, *Introduction to Quantum Groups*, Birkhäuser Boston, 1993.

[M] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS Conf. Math. Publ., **82**, Amer. Math. Soc., Providence, 1993.

[Ro] M. Rosso, *Lyndon bases and the multiplicative formula for R-matrices*, (2002), preprint.

Appendix: the proof of Lemma 3.6

Proof. (1): The proof is divided into 3 steps:

(i) If $j > i+3$, then taking decomposition $\mathcal{E}_{i,j'} = [\,\mathcal{E}_{i,i+2}, \mathcal{E}_{i+3,j'}]_{\bullet}$, combining with $\mathcal{E}_{i,i+2}e_{i+1} = e_{i+1}\mathcal{E}_{i,i+2}$ (by Lemma 3.3 (1)), $\mathcal{E}_{i+3,j'}e_{i+1} = e_{i+1}\mathcal{E}_{i+3,j'}$ (by Lemma 3.1 (2)), we get $\mathcal{E}_{i,j'}e_{i+1} = e_{i+1}\mathcal{E}_{i,j'}$.

(ii) If $j = i+3 = n$, then $\mathcal{E}_{n-3,n'}e_{n-2} = e_{n-2}\mathcal{E}_{n-3,n'}$ (by Lemma 3.3 (6)). If $j = i+3 < n$, then the decomposition $\mathcal{E}_{i,i+3'} = [\,\mathcal{E}_{i,i+4'}, e_{i+3}]_{\bullet}$ shows $\mathcal{E}_{i,i+3'}e_{i+1}$ $e_{i+1}\mathcal{E}_{i,i+3'}$, as e_{i+1} commutes with e_{i+3} and $\mathcal{E}_{i,i+4'} = [\,\mathcal{E}_{i,i+2}, \mathcal{E}_{i+3,i+4'}]_{\bullet}$ (by Lemma 3.1 (4), Lemma 3.3 (1) & Lemma 3.1 (2)).

(iii) If $j = i+1 = n$ (as $j \neq i+2$ by assumption), then the case is just (2.7). If $j = i+1 = n-1$, then by (2.4), Lemma 3.3 (1) & $(B5)$, we have

$$\begin{aligned}
[\,\mathcal{E}_{n-2,n-1'}, e_{n-1}] &= \mathcal{E}_{n-2,n-1'}e_{n-1} - r^{-2}e_{n-1}\mathcal{E}_{n-2,n-1'}\\
&= (\mathcal{E}_{n-2,n'}e_{n-1} - s^{-2}e_{n-1}\mathcal{E}_{n-2,n'})e_{n-1} - r^{-2}e_{n-1}(\mathcal{E}_{n-2,n'}e_{n-1} - s^{-2}e_{n-1}\mathcal{E}_{n-2,n'})\\
&= (\mathcal{E}_{n-2,n}e_n - rse_n\mathcal{E}_{n-2,n})e_{n-1}^2 - s^{-2}e_{n-1}(\mathcal{E}_{n-2,n}e_n - rse_n\mathcal{E}_{n-2,n})e_{n-1}\\
&\quad - r^{-2}e_{n-1}(\mathcal{E}_{n-2,n}e_n - rse_n\mathcal{E}_{n-2,n})e_{n-1} + (rs)^{-2}e_{n-1}^2(\mathcal{E}_{n-2,n}e_n - rse_n\mathcal{E}_{n-2,n})\\
&= \mathcal{E}_{n-2,n}\Big(e_ne_{n-1}^2 - (r^{-2}+s^{-2})e_{n-1}e_ne_{n-1} + (rs)^{-2}e_{n-1}^2e_n\Big)\\
&\quad -(rs)\Big(e_ne_{n-1}^2 - (r^{-2}+s^{-2})e_{n-1}e_ne_{n-1} + (rs)^{-2}e_{n-1}^2e_n\Big)\mathcal{E}_{n-2,n}\\
&= 0.
\end{aligned}$$

If $j = i+1 < n-1$, then by (ii), we have

$$\begin{aligned}
[\,\mathcal{E}_{i,i+1'}, e_{i+1}] &= \mathcal{E}_{i,i+1'}e_{i+1} - r^{-2}e_{i+1}\mathcal{E}_{i,i+1'}\\
&= (\mathcal{E}_{i,i+2'}e_{i+1} - s^{-2}e_{i+1}\mathcal{E}_{i,i+2'})e_{i+1} - r^{-2}e_{i+1}(\mathcal{E}_{i,i+2'}e_{i+1} - s^{-2}e_{i+1}\mathcal{E}_{i,i+2'})\\
&= (\mathcal{E}_{i,i+3'}e_{i+2} - s^{-2}e_{i+2}\mathcal{E}_{i,i+3'})e_{i+1}^2 - s^{-2}e_{i+1}(\mathcal{E}_{i,i+3'}e_{i+2} - s^{-2}e_{i+2}\mathcal{E}_{i,i+3'})e_{i+1}\\
&\quad -r^{-2}e_{i+1}(\mathcal{E}_{i,i+3'}e_{i+2} - s^{-2}e_{i+2}\mathcal{E}_{i,i+3'})e_{i+1} + (rs)^{-2}e_{i+1}^2(\mathcal{E}_{i,i+3'}e_{i+2} - s^{-2}e_{i+2}\mathcal{E}_{i,i+3'})\\
&= \mathcal{E}_{i,i+3'}\Big(e_{i+2}e_{i+1}^2 - (r^{-2}+s^{-2})e_{i+1}e_{i+2}e_{i+1} + (rs)^{-2}e_{i+1}^2e_{i+2}\Big)\\
&\quad -s^{-2}\Big(e_{i+2}e_{i+1}^2 - (r^{-2}+s^{-2})e_{i+1}e_{i+2}e_{i+1} + (rs)^{-2}e_{i+1}^2e_{i+2}\Big)\mathcal{E}_{i,i+3'}\\
&= 0,
\end{aligned}$$

where in the 1st " = " we used the definition, in the 2nd and 3rd " = " we used (2.4), in the 4th " = " we used $e_{i+1}\mathcal{E}_{i,i+3'} = \mathcal{E}_{i,i+3'}e_{i+1}$ in (ii), while in the last " = ", we used the (r,s)-Serre relations $(B5)$.

(2): From the proof of step (iii) in (1), we have $\mathcal{E}_{j-1,j'}e_j = r^{-2}e_j\mathcal{E}_{j-1,j'}$ for $j < n$. Hence, for $i < j-1$, we have $\mathcal{E}_{i,j'}e_j = r^{-2}e_j\mathcal{E}_{i,j'}$ since $\mathcal{E}_{i,j'} = [\,\mathcal{E}_{i,j-2}, \mathcal{E}_{j-1,j'}]_{r^2}$.

(3): If $i+2 = n$, then this is the special case of Lemma 3.4 (4).

If $i+2<n$, then using (2), we obtain

$$\begin{aligned}
[\,\mathcal{E}_{i,i+1'}, e_{i+2}] &= \mathcal{E}_{i,i+1'}e_{i+2} - e_{i+2}\mathcal{E}_{i,i+1'}\\
&= \Big(\mathcal{E}_{i,i+2'}e_{i+1} - s^{-2}e_{i+1}\mathcal{E}_{i,i+2'}\Big)e_{i+2} - e_{i+2}\Big(\mathcal{E}_{i,i+2'}e_{i+1} - s^{-2}e_{i+1}\mathcal{E}_{i,i+2'}\Big)\\
&= \mathcal{E}_{i,i+2'}\Big(e_{i+1}e_{i+2} - r^2 e_{i+2}e_{i+1}\Big) - (rs)^{-2}\Big(e_{i+1}e_{i+2} - r^2e_{i+2}e_{i+1}\Big)\mathcal{E}_{i,i+2'}\\
&= \mathcal{E}_{i,i+2'}\mathcal{E}_{i+1,i+2} - (rs)^{-2}\mathcal{E}_{i+1,i+2}\mathcal{E}_{i,i+2'},
\end{aligned}$$

so (3) is equivalent to prove

$$\mathcal{E}_{i+1,i+2}\mathcal{E}_{i,i+2'} = (rs)^2\mathcal{E}_{i,i+2'}\mathcal{E}_{i+1,i+2}, \qquad i+2<n.$$

If $i+2=n-1$, then using Lemmas 3.4 (1) and 3.2 (3), we obtain

$$\begin{aligned}
\mathcal{E}_{n-2,n-1}\mathcal{E}_{n-3,n-1'} &= \mathcal{E}_{n-2,n-1}\mathcal{E}_{n-3,n'}e_{n-1} - s^{-2}\mathcal{E}_{n-2,n-1}e_{n-1}\mathcal{E}_{n-3,n'}\\
&= \Big(r^2\mathcal{E}_{n-3,n'}\mathcal{E}_{n-2,n-1} + \mathcal{E}_{n-3,n}\mathcal{E}_{n-2,n} - rs\mathcal{E}_{n-2,n}\mathcal{E}_{n-3,n}\Big)e_{n-1}\\
&\quad - e_{n-1}\mathcal{E}_{n-2,n-1}\mathcal{E}_{n-3,n'}\\
&= (rs)^2\mathcal{E}_{n-3,n'}e_{n-1}\mathcal{E}_{n-2,n-1}\\
&\quad + \underline{\mathcal{E}_{n-3,n}\mathcal{E}_{n-2,n}e_{n-1} - rs\mathcal{E}_{n-2,n}\mathcal{E}_{n-3,n}e_{n-1}}\\
&\quad - r^2e_{n-1}\mathcal{E}_{n-3,n'}\mathcal{E}_{n-2,n-1}\\
&\quad - \underline{e_{n-1}\mathcal{E}_{n-3,n}\mathcal{E}_{n-2,n} + rse_{n-1}\mathcal{E}_{n-2,n}\mathcal{E}_{n-3,n}}\\
&= (rs)^2\mathcal{E}_{n-3,n-1'}\mathcal{E}_{n-2,n-1},
\end{aligned}$$

where the sum of terms underlined is 0 since e_{n-1} commutes with $\mathcal{E}_{n-2,n}, \mathcal{E}_{n-3,n}$. The argument for $i+2<n-1$ is similar by using Lemmas 3.3 (8) and 3.4 (2).

(4) For $i<j<l<n$: first noting the case when $j=l-1$ and $i=j-1=l-2$, by (3), we have $\mathcal{E}_{l-2,l-1'}e_l = e_l\mathcal{E}_{l-2,l-1'}$ for $l<n$. For the case when $j=l-1$ and $i\le l-3$, using $\mathcal{E}_{i,l-1'} = [\,\mathcal{E}_{i,l-3}, \mathcal{E}_{l-2,l-1'}]_{r^2}$ and $e_l\mathcal{E}_{i,l-3} = \mathcal{E}_{i,l-3}e_l$, we obtain $e_l\mathcal{E}_{i,l-1'} = \mathcal{E}_{i,l-1'}e_l$.

For the case when $j<l-1$, noting that $\mathcal{E}_{i,j'} = [\cdots[\,\mathcal{E}_{i,l-1'}, e_{l-2}]_{s^{-2}}, \cdots, e_j]_{s^{-2}}$ and $e_le_k = e_ke_l$ for $j\le k\le l-2$, we get $e_l\mathcal{E}_{i,j'} = \mathcal{E}_{i,j'}e_l$ for $i<j<l<n$.

(5): First, we claim: $\mathcal{E}_{i,n'}\mathcal{E}_{i,n-1'} = r^{-2}\mathcal{E}_{i,n-1'}\mathcal{E}_{i,n'}$ for $i<n-1$. Indeed, by Lemma 3.4 (1), $e_{n-1}\mathcal{E}_{i,n'} = r^2\mathcal{E}_{i,n'}e_{n-1} + \mathcal{E}_{i,n}\mathcal{E}_{n-1,n} - rs\mathcal{E}_{n-1,n}\mathcal{E}_{i,n}$. Again by Lemma 3.5 (3) & Lemma 3.4 (5), $\mathcal{E}_{i,n}\mathcal{E}_{i,n'} = s^2\mathcal{E}_{i,n'}\mathcal{E}_{i,n}$, $\mathcal{E}_{i,n'}\mathcal{E}_{n-1,n} = \mathcal{E}_{n-1,n}\mathcal{E}_{i,n'}$. So using (2.4), we have

$$\begin{aligned}
[\,\mathcal{E}_{i,n'}, \mathcal{E}_{i,n-1'}] &= \mathcal{E}_{i,n'}\mathcal{E}_{i,n-1'} - r^{-2}\mathcal{E}_{i,n-1'}\mathcal{E}_{i,n'}\\
&= \mathcal{E}_{i,n'}^2e_{n-1} - (s^{-2}+r^{-2})\mathcal{E}_{i,n'}e_{n-1}\mathcal{E}_{i,n'} + (rs)^{-2}(e_{n-1}\mathcal{E}_{i,n'})\mathcal{E}_{i,n'}\\
&= \mathcal{E}_{i,n'}^2e_{n-1} - r^{-2}\mathcal{E}_{i,n'}(e_{n-1}\mathcal{E}_{i,n'}) + (rs)^{-2}(\mathcal{E}_{i,n}\mathcal{E}_{n-1,n} - rs\mathcal{E}_{n-1,n}\mathcal{E}_{i,n})\mathcal{E}_{i,n'}\\
&= -r^{-2}\mathcal{E}_{i,n'}\Big(\mathcal{E}_{i,n}\mathcal{E}_{n-1,n} - rs\mathcal{E}_{n-1,n}\mathcal{E}_{i,n}\Big) + (rs)^{-2}\Big(\mathcal{E}_{i,n}\mathcal{E}_{n-1,n} - rs\mathcal{E}_{n-1,n}\mathcal{E}_{i,n}\Big)\mathcal{E}_{i,n'}\\
&= 0,
\end{aligned}$$

this means $\mathcal{E}_{i,n'}\mathcal{E}_{i,n-1'} = r^{-2}\mathcal{E}_{i,n-1'}\mathcal{E}_{i,n'}$ for $i<n-1$.

Next, we claim: $\mathcal{E}_{i,n'}\mathcal{E}_{i,j'} = r^{-2}\mathcal{E}_{i,j'}\mathcal{E}_{i,n'}$ for $i<j<n$. In fact, noting $\mathcal{E}_{i,j'} = [\cdots[\,\mathcal{E}_{i,n-1'}, e_{n-2}]_{s^{-2}}, \cdots, e_j]_{s^{-2}}$ and $e_k\mathcal{E}_{i,n'} = \mathcal{E}_{i,n'}e_k$ for any k with $i<k<n-1$, we arrive at the required result.

Finally, we claim: $\mathcal{E}_{i,l'}\mathcal{E}_{i,j'} = r^{-2}\mathcal{E}_{i,j'}\mathcal{E}_{i,l'}$ for $j < l < n$. This follows from (4) and $\mathcal{E}_{i,l'} = [\cdots[\,\mathcal{E}_{i,n'}, e_{n-1}]_{s^{-2}}, \cdots, e_l]_{s^{-2}}$.

(6) follows from (4) and Lemma 3.5 (6).

(7) follows from (2) & (6).

(8): By Lemma 3.3 (5), we have $\mathcal{E}_{i,n-1}\mathcal{E}_{i,n} = s^2\mathcal{E}_{i,n}\mathcal{E}_{i,n-1}$ and

$$\begin{aligned}
&\mathcal{E}_{i,n-1}\mathcal{E}_{i,n'}-(rs)^2\mathcal{E}_{i,n'}\mathcal{E}_{i,n-1}\\
&\quad= \mathcal{E}_{i,n-1}(\mathcal{E}_{i,n}e_n-rse_n\mathcal{E}_{i,n})-(rs)^2(\mathcal{E}_{i,n}e_n-rse_n\mathcal{E}_{i,n})\mathcal{E}_{i,n-1}\\
&\quad= s^2\mathcal{E}_{i,n}(\mathcal{E}_{i,n-1}e_n)-rs(\mathcal{E}_{i,n-1}e_n)\mathcal{E}_{i,n}-(rs)^2\mathcal{E}_{i,n}(e_n\mathcal{E}_{i,n-1})+r^3s(e_n\mathcal{E}_{i,n-1})\mathcal{E}_{i,n}\\
&\quad= s(s-r)\mathcal{E}_{i,n}^2.
\end{aligned}$$

This completes the proof. □

Department of Mathematics, East China Normal University, Min Hang Campus, Dong Chuan Road 500, Shanghai 200241, PR China
E-mail address: `nhhu@math.ecnu.edu.cn`

School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, PR China
E-mail address: `xiulingwang@nankai.edu.cn`